



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>6</sup> : <b>C07D 453/02, A61K 31/435, C07D 451/02, 471/08, 487/08, 211/62, A61K 31/445</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 99/21855</b></p> <p>(43) International Publication Date: 6 May 1999 (06.05.99)</p>
<p>(21) International Application Number: PCT/GB98/03211</p> <p>(22) International Filing Date: 27 October 1998 (27.10.98)</p> <p>(30) Priority Data: 9722817.5 29 October 1997 (29.10.97) GB 9813689.8 25 June 1998 (25.06.98) GB</p> <p>(71) Applicants (for all designated States except US): SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, P.O. Box 7929, Philadelphia, PA 19103 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): BERRY, Valerie [GB/US]; SmithKline Beecham Pharmaceuticals, 1250 South Collegeville Road, P.O. Box 5089, Collegeville, PA 19426-0989 (US). DABBS, Stephen [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). FRYDRYCH, Colin, Henry [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). HUNT, Eric [GB/GB]; SmithKline Beecham Pharmaceuticals, New</p>		<p>Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). WOODNUTT, Gary [GB/US]; SmithKline Beecham Pharmaceuticals, 1250 South Collegeville Road, P.O. Box 5089, Collegeville, PA 19426-0989 (US). SANDERSON, Francis, Dominic [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).</p> <p>(74) Agent: CONNELL, Anthony, Christopher; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
<p>(54) Title: PLEUROMUTILIN DERIVATIVES AS ANTIMICROBIALS</p>		
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <p>(IA)</p> </div> <div style="text-align: center;"> <p>(IB)</p> </div> </div>		
<p>(57) Abstract</p> <p>Derivatives of pleuromutilin of formula (IA) or (IB) in which the glycolic ester moiety at position (14) is replaced by <math>R^2(CH_2)_mX(CH_2)_nCH_2COO</math> are of use in antimicrobial therapy. In which: each of n and m is independently 0, 1 or 2; X is selected from -O-, -S-, S(O)-, -SO<sub>2</sub>-, -CO.O-, -NH-, -CONH-, NHCONH- and a bond; R<sup>1</sup> is vinyl or ethyl; R<sup>2</sup> is a non-aromatic monocyclic or bicyclic group containing one or two basic nitrogen atoms and attached through a ring carbon atom; R<sup>3</sup> is H or OH; or the moiety <math>R^2(CH_2)_mX(CH_2)_nCH_2COO</math> at position 14 of (IA) or (IB) is replaced by <math>R^aR^bC=CHCOO</math> in which one of R<sup>a</sup> and R<sup>b</sup> is hydrogen and the other is R<sup>2</sup> or R<sup>a</sup> and R<sup>b</sup> together form R<sup>2</sup>; or a pharmaceutically acceptable salt thereof.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**



Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## PLEUROMUTILIN DERIVATIVES AS ANTIMICROBIALS

The present invention relates to novel compounds, to processes for their preparation, to pharmaceutical compositions containing them and to their use in medical therapy, particularly antibacterial therapy.

Pleuromutilin, the compound of formula (A), is a naturally occurring antibiotic which has antimycoplasmal activity and modest antibacterial activity. It has been shown that the antimicrobial activity can be improved by replacing the glycolic ester moiety at position 14 by an R-X-CH<sub>2</sub>CO<sub>2</sub>- group, where R is an aliphatic or aromatic moiety and X is O, S, or NR' (H Egger and H Reinshagen, J Antibiotics, 1976, 29, 923). Tiamulin, the compound of formula (B), which is used as a veterinary antibiotic, is a derivative of this type (G Hogenauer in Antibiotics, Vol. V, part 1, ed. F E Hahn, Springer-Verlag, 1979, p.344).



In this application, the non-conventional numbering system which is generally used in the literature (G Hogenauer, loc.cit.) is used.

WO 97/25309 (SmithKline Beecham) describes further modification of the acyloxy group, disclosing 14-O-carbamoyl derivatives of mutilin or 19, 20-dihydromutilin, in which the N-atom of the carbamoyl group is unsubstituted, mono- or di-substituted.



WO 98/05659 (SmithKline Beecham) discloses 14-O-carbamoyl derivatives of mutilin or 19, 20-dihydromutilin, in which the N-atom of the carbamoyl group is acylated by a group which includes an azabicyclic moiety.

WO 98/14189 (SmithKline Beecham, International Publication Date 9 April 1998) discloses the use of the topical antibacterial agent mupirocin for treating bacterial infections associated with the colonisation of the nasopharynx by pathogenic organisms, in particular, the prophylactic treatment of recurrent sinusitis and recurrent otitis media, especially with novel spray or cream formulations adapted for administration to the

nasopharynx. In addition, Nsouli (Annals of Allergy, Asthma and Immunology, January 1996, 76(1), 117) has described a clinical study involving the use of a 0.2% aqueous solution of mupirocin in reducing the attacks of sinusitis.

- 5 We have now found that further novel pleuromutilin derivatives have improved antimicrobial properties.

Accordingly, the present invention provides a compound of general formula (IA) or (IB):



in which:

- 15 each of n and m is independently 0, 1 or 2;  
 X is selected from -O-, -S-, -S(O)-, -SO₂-, -CO.O-, -NH-, -CONH-, -NHCONH- and a bond;  
 R¹ is vinyl or ethyl;  
 R² is a non-aromatic monocyclic or bicyclic group containing one or two basic nitrogen  
 20 atoms and attached through a ring carbon atom;  
 R³ is H or OH; or  
 the moiety R²(CH₂)ₘX(CH₂)ₙCH₂COO at position 14 of (IA) or (IB) is replaced by  
 RᵃRᵇC=CHCOO in which one of Rᵃ and Rᵇ is hydrogen and the other is R² or Rᵃ and Rᵇ  
 together form R², or  
 25 a pharmaceutically acceptable salt thereof.

When R<sup>2</sup> is monocyclic, it typically contains from 4 to 8 ring atoms, and, when bicyclic, it typically contains from 5 to 10 ring atoms in each ring, and is optionally substituted on carbon by up to 3 substituents. Suitable substituents include alkyl, alkyloxy, alkenyl and alkenyloxy, each of which may be carried by either a bridgehead or a non-bridgehead carbon atom. In addition, the or each nitrogen atom may be substituted by oxygen, to form an N-oxide, or by mono- or dialkyl, in which case it will be appreciated that a quaternary cation can be formed. The counterion may be a halide ion such as chloride or bromide, preferably chloride. The aza ring system additionally may contain one or more double bonds.

10

Representative bicyclic and monocyclic groups for R<sup>2</sup> include piperidinyl, pyrrolidyl, quinuclidinyl, azabicyclo[2.2.1]heptyl, azabicyclo[4.3.0]nonyl, azabicyclo[3.2.1]octyl, azabicyclo[3.3.0]octyl, azabicyclo[2.2.2]octyl, azabicyclo[3.2.1]octenyl, azabicyclo[3.3.1]nonyl and azabicyclo[4.4.0]decyl, all of which may be substituted or unsubstituted. Preferred examples for R<sup>2</sup> include quinuclidinyl.

15

The compounds of formula (IA) in which R<sup>3</sup> is hydroxy have the (2S) configuration at the carbon bearing this hydroxy group.

20 Preferably, n is 0. Preferably, m is 0 or 1.

Preferred compounds are those of formula (IA).

Alkyl and alkenyl groups referred to herein include straight and branched groups containing up to six carbon atoms and are optionally substituted by one or more groups selected from the group consisting of aryl, heterocyclyl, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)alkylthio, aryl(C<sub>1-6</sub>)alkoxy, aryl(C<sub>1-6</sub>)alkylthio, amino, mono- or di-(C<sub>1-6</sub>)alkylamino, cycloalkyl, cycloalkenyl, carboxy and esters thereof, amides of carboxy, ureido, carbamimidoyl (amidino), guanidino, alkyl-sulfonyl, amino-sulfonyl (C<sub>1-6</sub>)acyloxy, (C<sub>1-6</sub>)acylamino, azido, hydroxy, and halogen.

25

30

Cycloalkyl and cycloalkenyl groups referred to herein include groups having from three to eight ring carbon atoms and are optionally substituted as described hereinabove for alkyl and alkenyl groups.

35

When used herein, the term "aryl" means single and fused rings suitably containing from 4 to 7, preferably 5 or 6, ring atoms in each ring, which rings may each be unsubstituted or substituted by, for example, up to three substituents. A fused ring system may include

aliphatic rings and need include only one aromatic ring. Representative aryl groups include phenyl and naphthyl such as 1-naphthyl or 2-naphthyl.

- Suitably any aryl group, including phenyl and naphthyl, may be optionally substituted by up to five, preferably up to three substituents. Suitable substituents include halogen, (C<sub>1-6</sub>)alkyl, aryl, aryl(C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)alkoxy(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkyl, aryl(C<sub>1-6</sub>)alkoxy, hydroxy, nitro, cyano, azido, amino, mono- and di-*N*-(C<sub>1-6</sub>)alkylamino, acylamino, arylcarbonylamino, acyloxy, carboxy, carboxy salts, carboxy esters, carbamoyl, mono- and di-*N*-(C<sub>1-6</sub>)alkylcarbamoyl, (C<sub>1-6</sub>)alkoxycarbonyl, aryloxycarbonyl, ureido, guanidino, sulphonylamino, aminosulphonyl, (C<sub>1-6</sub>)alkylthio, (C<sub>1-6</sub>)alkyl sulphinyl, (C<sub>1-6</sub>)alkylsulphonyl, heterocyclyl and heterocyclyl (C<sub>1-6</sub>)alkyl. In addition, two adjacent ring carbon atoms may be linked by a (C<sub>3-5</sub>)alkylene chain, to form a carbocyclic ring.
- When used herein the terms "heterocyclyl" and "heterocyclic" suitably include, unless otherwise defined, aromatic and non-aromatic, single and fused, rings suitably containing up to four heteroatoms in each ring, each of which is selected from oxygen, nitrogen and sulphur, which rings, may be unsubstituted or substituted by, for example, up to three substituents. Each heterocyclic ring suitably has from 4 to 7, preferably 5 or 6, ring atoms.
- A fused heterocyclic ring system may include carbocyclic rings and need include only one heterocyclic ring.

- Preferably substituents for a heterocyclyl group are selected from halogen, (C<sub>1-6</sub>)alkyl, aryl(C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)alkoxy(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkyl, hydroxy, amino, mono- and di-*N*-(C<sub>1-6</sub>)alkyl-amino, acylamino, carboxy, carboxy salts, carboxy esters, carbamoyl, mono- and di-*N*-(C<sub>1-6</sub>)alkylcarbonyl, aryloxycarbonyl, (C<sub>1-6</sub>)alkoxycarbonyl(C<sub>1-6</sub>)alkyl, aryl, oxy groups, ureido, guanidino, sulphonylamino, aminosulphonyl, (C<sub>1-6</sub>)alkylthio, (C<sub>1-6</sub>)alkylsulphinyl, (C<sub>1-6</sub>)alkylsulphonyl, heterocyclyl and heterocyclyl(C<sub>1-6</sub>)alkyl.

- Depending on the position of attachment of substituents, two or more diastereoisomers may be possible. In that situation the present invention includes the individual diastereoisomers and mixtures thereof.

- Preferred examples of compounds of the invention include:
- Mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate;
- Mutilin 14-(quinuclid-4-ylmethylsulfanyl)-acetate;
- Mutilin-14-(1-methylpiperid-4-ylsulfanyl)-acetate; and
- Mutilin 14-(*exo*-8-methyl-8-azabicyclo[3.2.1]oct-3-ylsulfanyl)-acetate.

The compounds of this invention may be in crystalline or non-crystalline form, and, if crystalline, may optionally be solvated, especially hydrated. This invention includes within its scope stoichiometric hydrates as well as compounds containing variable amounts of water.


The compounds according to the invention are suitably provided in substantially pure form, for example at least 50% pure, suitable at least 60% pure, advantageously at least 75% pure, preferably at least 85% pure, more preferably at least 95% pure, especially at least 98% pure, all percentages being calculated as weight/weight.

The compounds of the invention may be in the form of free bases or acid addition salts. Compounds carrying a carboxy substituent may be in the form of zwitterions, or alkali metal salts (of the carboxy group). Pharmaceutically acceptable salts are preferred.


Pharmaceutically acceptable acid-addition salts include those described by Berge, Bighley, and Monkhouse, *J. Pharm. Sci.*, 1977, **66**, 1-19. Suitable salts include the hydrochloride, maleate, and methanesulphonate; particularly the hydrochloride.

Compounds of the present invention may be readily prepared from available starting materials by adapting synthetic processes well known in the art.

Accordingly, in a first aspect, the present invention provides a process for preparing a compound of formula (I) which comprises reacting a compound of formula (IIA) or (IIB):




(IIA)



(IIB)

in which Y is hydrogen or a removable hydroxy-protecting group, and R<sup>1A</sup> and R<sup>3A</sup> are R<sup>1</sup> and R<sup>3</sup> are as defined for formulae (IA) and (IB) or groups convertible to R<sup>1</sup> and R<sup>3</sup>,

with an active derivative of a carboxylic acid of formula (III):



(III)

where  $R^{2A}$  is  $R^2$  as defined for formulae IA and IB or a group convertible to  $R^2$ , under ester forming conditions and, where required or desired,

- 5 converting Y to hydrogen,
- converting an  $R^{1A}$ ,  $R^{2A}$  or  $R^{3A}$  group to a  $R^1$ ,  $R^2$  or  $R^3$  group, and/or
- converting one  $R^1$ ,  $R^2$  or  $R^3$  group to another  $R^1$ ,  $R^2$  or  $R^3$  group.

- 10 Conventional methods for ester formation are described in the literature, for example in *Comprehensive Organic Functional Group Transformations*, Vol. 5, ed. C J Moody, p. 123-130, Elsevier Scientific, Oxford, 1995. The active derivative used as an acylating agent may be for example an acid chloride, acid bromide, a mixed anhydride, or an *N*-acyl-imidazole. The preferred agent is an acid chloride. General methods for forming such acylating agents are described in the chemical literature (see I O Sutherland,
- 15 *Comprehensive Organic Chemistry*, Vol. 2, ed. I O Sutherland, pages 875-883 (Pergamon Press, Oxford, 1979), and references therein).

- The ester-forming reaction can be carried out in the presence of an organic base, an inorganic base, or an acid. Organic bases include pyridine, 2,6-lutidine, triethylamine, and
- 20 *N,N*-dimethylaniline. Inorganic bases include sodium hydride, lithium hydride, potassium carbonate, lithium hexamethyldisilazide, and sodium hexamethyldisilazide. Acids include *p*-toluenesulphonic acid, benzene sulphonic acid, and sulphuric acid. Optionally, when the reaction is carried out in the presence of a base, an acylation catalyst (G Hofle and W Steglich, *Synthesis*, 1972, 619) such as 4-dimethylamino-pyridine or 4-pyrrolidino-
- 25 pyridine may also be added to the reaction mixture. Solvents for the ester forming reaction include tetrahydrofuran, 1,4-dioxane, acetonitrile, *N,N*-dimethylformamide, diethyl ether, dichloromethane, and chloroform. A preferred solvent is tetrahydrofuran.

- Useful methods for acylating the 14-hydroxyl in the present invention include the use of
- 30 the following: acid chloride in *N,N*-dimethylformamide at elevated temperature (*e.g.* 100°C to 120°C); acid chloride in the presence of an organic base (*e.g.* pyridine, 2,6-lutidine, 2,4,6-collidine, di-*iso*-propylethylamine) or an inorganic base (*e.g.* sodium or lithium hexamethyldisilazide); carboxylic acid in the presence of dicyclohexylcarbodiimide and an acylation catalyst (*e.g.* 4-dimethylamino-pyridine, 4-pyrrolidino-pyridine); a mutilin
- 35 14-chloroformate derivative plus carboxylic acid, tertiary base (*e.g.* triethylamine, di-*iso*-propyl-ethylamine), and an acylation catalyst (*e.g.* 4-dimethylamino-pyridine, 4-pyrrolidino-pyridine).



Conversions of an  $R^{1A}$ ,  $R^{2A}$  or  $R^{3A}$  group to a  $R^1$ ,  $R^2$  or  $R^3$  group typically arise when a protecting group is needed during the above coupling reaction or during the preparation of the reactants by the procedures described below. Interconversion of one  $R^1$ ,  $R^2$  or  $R^3$  group to another typically arises when one compound of formula IA/B is used as the  
5 immediate precursor of another compound of formula IA/B or when it is easier to introduce a more complex or reactive substituent at the end of a synthetic sequence.

Preferably Y is a hydroxyl protecting group such as an acyl group, for example so that -OY is trifluoroacetyl or dichloroacetyl. When the intended  $R^3$  is also hydroxyl, then  $R^{3A}$  is  
10 also preferably acyloxy, for example acetyl or dichloroacetyl. Hydroxyl groups at positions 11 and 2 (as groups OY and  $R^{3A}$ ) may be protected using, for example, dichloroacetic anhydride and pyridine in tetrahydrofuran or *N*-trifluoroacetyl-imidazole in tetrahydrofuran at 0°C. After the reaction with the derivative of acid III is complete the protecting acyl groups may be removed to restore the hydroxyl groups by hydrolysis *e.g.* using NaOH in  
15 MeOH.


It may also be necessary to protect substituent groups in the acid component (III) prior to reaction with the compound of formulae (IIA) or (IIB), for example protecting N atoms with alkoxycarbonyl, for example *t*-butoxycarbonyl.

20 Suitable hydroxy, carboxy and amino protecting groups are those well known in the art and which may be removed under conventional conditions and without disrupting the remainder of the molecule. A comprehensive discussion of the ways in which hydroxy, carboxy and amino groups may be protected and methods for cleaving the resulting  
25 protected derivatives is given in for example "Protective Groups in Organic Chemistry" (T.W. Greene, Wiley-Interscience, New York, 2nd edition, 1991). Particularly suitable hydroxy protecting groups include, for example, triorganosilyl groups such as, for instance, trialkylsilyl and also organocarbonyl and organooxycarbonyl groups such as, for instance, acetyl, allyloxycarbonyl, 4-methoxybenzyloxycarbonyl and 4-nitrobenzyloxycarbonyl.  
30 Particularly suitable carboxy protecting groups include alkyl and aryl groups, for instance methyl, ethyl and phenyl. Particularly suitable amino protecting groups include alkoxycarbonyl, 4-methoxybenzyloxycarbonyl and 4-nitrobenzyloxycarbonyl.

$R^{1A}$  is typically the  $R^1$  group vinyl, and this may be converted to the alternative  $R^1$  ethyl group by hydrogenating the vinyl group to form an ethyl group, typically by hydrogenation over a palladium catalyst (*e.g.* 10% palladium-on-carbon) in a solvent such as ethyl acetate, ethanol, dioxane, or tetrahydrofuran.

$R^{3A}$  is typically hydrogen or protected hydroxyl, such as acyloxy. After the coupling reaction, protecting acyl groups may be removed to restore the hydroxyl groups by hydrolysis *e.g.* using NaOH in MeOH.

- 5 Alternatively a compound of formula (IA) in which  $R^3$  is hydrogen may be prepared by treating a compound of formula (IIC):




(IIC)

- 10 where  $R^{1A}$  is as defined for formulae (IIA) and (IIB),  
with an active derivative of the acid of formula (III) under ester forming conditions, and  
then treating the product with an acid, and, where required or desired,  
converting an  $R^{1A}$  or  $R^{2A}$  group to a  $R^1$  or  $R^2$  group, and/or  
converting one  $R^1$  or  $R^2$  group to another  $R^1$  or  $R^2$  group.
- 15 The acid treatment indicated above converts the epi-mutilin configuration of formula (IIC)  
to the usual mutilin nucleus of formula (IIA). Typically this conversion is carried out by  
treatment with conc. HCl or Lukas reagent (conc. HCl saturated with  $ZnCl_2$ ) in dioxane.
- 20 As in formulae (IIA) and (IIB),  $R^{2A}$  is typically the  $R^2$  group vinyl, and this may be  
converted to the alternative  $R^2$  group by hydrogenating the vinyl group to form an ethyl  
group. Also it may again be necessary to protect substituent groups in the derivative of  
acid component (III) prior to reaction, for example protecting N atoms with, for example,  
*t*-butoxycarbonyl


- 25 In cases where the intermediate of formula (IIA) and (IIB) (such as Y= acetyl) are used, a  
base-labile protecting group may conveniently be removed at the same time as the group Y  
is deprotected. In cases when the intermediate of formula (IIC) is used, an acid-labile  
protecting group may conveniently be removed at the same time as the acid treatment that  
30 converts the epi-mutilin configuration into the desired configuration of the compounds of  
the invention.

The compounds of formulae (IIA), (IIB) and (IIC) may be prepared from compounds of  
formulae (IV) and (V)

35



(IV)



(V)

Suitable compounds as formula (IV) include 11-O-acyl mutilin derivatives, e.g. mutilin 11-acetate (A J Birch, C W Holzapfel, R W Richards, *Tetrahedron (Suppl.)*, 1966, 8, Part II, 359) or mutilin 11-dichloroacetate or mutilin 11-trifluoroacetate. Formula (V) is (3R)-3-deoxy-11-deoxy-3-methoxy-11-oxo-4-epi-mutilin (H Berner, G Schulz and H Schneider, *Tetrahedron*, 1980, 36, 1807).


- Compounds (IV) and (V) are effectively the compounds of formula (IIA) and (IIC) respectively in which  $R^{1A}$  is vinyl and  $R^{3A}$  is hydrogen (compound IIA). They may be converted into the corresponding compounds in which  $R^{1A}$  is ethyl by hydrogenation, typically by hydrogenation over a palladium catalyst (e.g. 10% palladium-on-carbon) in a solvent such as ethyl acetate, ethanol, dioxane, or tetrahydrofuran.
- Copmpounds of formula (IIA) in which  $R^{3A}$  is hydroxyl may be obtained by first preparing 2-hydroxymethylene mutilin from a compound of formula (IV). Using procedures based on that described by A.J. Birch, C.W. Holzapfel and R.W. Rickards (*Tet (Suppl)* 1996 8 part III 359), a compound of formula (IV) in toluene and methyl formate is treated with sodium methoxide and stirred under argon. The product is a mixture of the desired 2-hydroxymethylene compound and corresponding compounds substituted by formate at position 11 (if OY is OH) and /or position 14. The formate groups may be removed when desired by treatment with potassium hydroxide in methanol.

The product mixture may however be used directly to prepare 2-diazo-mutilin derivatives using the method described by H Berner, G Schulz, and G Fisher, *Monatsh. Chem.*, 1981, 112, 1441, for example reacting a solution of a 2-hydroxymethylene-mutilin and the formate derivatives in dichloromethane at -10 °C under argon with tosylazide and triethylamine. Removal of the formate groups as described above leaves 2-diazo-mutilin, which may be reacted with a carboxylic acid to give a 2-acyloxy-mutilin, effectively a compound of formula (IIA) in which  $R^{3A}$  is protected hydroxyl. Suitably reaction with dichloroacetic acid gives 2-dichloroacetoxy-mutilin, which can be deprotected as described above to provide 2-OH, preferably after coupling with the derivative of acid (III). This reaction produces (2S)-2-hydroxy derivatives.

Compounds of formula (IIB) are either 1,2-didehydro-mutilin or obtainable therefrom by manipulation of OY and R<sup>1A</sup> as described above. 1,2-Didehydro-mutilins may be prepared using the method described by G Schulz and H Berner in *Tetrahedron*, 1984, 40, 905.

The above described modifications to the mutilin nucleus may also be carried out after coupling of compounds of formula (IIA) and (IIC) where R<sup>3A</sup> is hydrogen (*i.e.* based on mutilin and epi-mutilin) with the active derivative of acid (III).

In another aspect, the present invention provides a method for preparing compounds of the invention in which X is O, S, NH, CO.O or CONH which comprises reacting a compound of formula VIA or VIB



where Y is hydrogen or a removable hydroxy-protecting group, and R<sup>1A</sup> and R<sup>3A</sup> are R<sup>1</sup> and R<sup>3</sup> as defined for formulae IA and IB or groups convertible to R<sup>1</sup> and R<sup>3</sup>, n is as defined for formulae IA and IB, and R<sup>L</sup> is a leaving group or OH or NH<sub>2</sub>, with a compound of formula (VII):



where R<sup>2A</sup> is R<sup>2</sup> as defined for formulae (IA) and (IB) or a group convertible to R<sup>2</sup>, and X and m are as defined for formulae IA and IB, or

when X is CO.O with an active derivative of the acid of formula (VII), by one of the procedures set out below,

and where required or desired

converting Y to hydrogen,

converting an R<sup>1A</sup>, R<sup>2A</sup> or R<sup>3A</sup> group to an R<sup>1</sup>, R<sup>2</sup> or R<sup>3</sup> group, and/or converting one R<sup>1</sup>, R<sup>2</sup> or R<sup>3</sup> group to another R<sup>1</sup>, R<sup>2</sup> or R<sup>3</sup> group.

As in the method discussed above starting from compounds (IIA/B/C), preferably Y is a hydroxyl protecting group such as an acyl group, for example so that -OY is trifluoroacetyl or dichloroacetyl. When the intended  $R^3$  is also hydroxyl then  $R^{3A}$  is also preferably acyloxy, for example acetyl or dichloroacetyl.

5

It may also be necessary to protect substituent groups in the compound of formula (VII) prior to reaction with the compound (VIA) or (VIB), for example protecting N atoms with alkoxycarbonyl, for example *t*-butoxycarbonyl.

10 Suitable hydroxy, carboxy and amino protecting groups are those well known in the art and are discussed above.

$R^{1A}$  is typically the  $R^1$  group vinyl, and this may be converted to the alternative  $R^1$  ethyl group by hydrogenating the vinyl group to form an ethyl group, typically by hydrogenation over a palladium catalyst (*e.g.* 10% palladium-on-carbon) in a solvent such as ethyl acetate, ethanol, dioxane, or tetrahydrofuran.

15

$R^{3A}$  is typically hydrogen or protected hydroxyl, such as acyloxy. After the coupling reaction, protecting acyl groups may be removed to restore the hydroxyl groups by hydrolysis *e.g.* using NaOH in MeOH.

20

Procedures for coupling the group  $R^L(CH_2)_nCH_2CO.O-$  with compound  $R^{2A}-(CH_2)_m-XH$  include the following:

25 (a) when  $R^L$  is a leaving group, such as 4-MeC<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>O, MeSO<sub>2</sub>O, F<sub>3</sub>CSO<sub>2</sub>O, Br or Cl, and X is O, S or NH:

(i) where X = O, the alcohol  $R^2-(CH_2)_m-OH$  may be converted into the alkoxide by reaction with an inorganic base, such as sodium hydride, lithium hydride, sodium hexamethyldisilazide, or lithium hexamethyldisilazide, in a non-hydroxylic solvent, such as *N,N*-dimethylformamide or tetrahydrofuran, prior to reaction with the compound of formula VIA/B;

30

(ii) where X = S, the thiol  $R^2-(CH_2)_m-SH$  may be reacted with the compound of formula VIA/B in the presence of an inorganic base, such as sodium methoxide, sodium ethoxide, sodium hydride, sodium hexamethyldisilazide, or lithium hexamethyldisilazide, in a solvent such as 2-propanol, ethanol, methanol, *N,N*-dimethylformamide, or tetrahydrofuran.

35

(iii) where X = NH, the amine  $R^2-(CH_2)_m-NH_2$  may be reacted with the compound of formula VIA/B in a solvent such as *N,N*-dimethylformamide or tetrahydrofuran, optionally


in the presence of a base such as potassium carbonate, pyridine, *N,N*-di-(*iso*-propyl)-ethylamine, or triethylamine.

- 5 (b) when X is CONH, a compound of formula VIA/B in which  $R^L$  is amino may be reacted with a compound of formula  $R^{2A}-(CH_2)_m-CO_2H$ , or an acylating agent derived therefrom, using one of the general methods for amide formation that are described in the chemical literature. General methods for amide formation are described by B C Challis and J A Challis in *Comprehensive Organic Chemistry*, Vol. 2, ed. I O Sutherland, pages 959-964 (Pergamon Press, Oxford, 1979).

10

- (c) when X is CO.O, a compound of formula VIA/B in which  $R^L$  is hydroxy may be reacted with an acylating agent derived from a compound of formula  $R^{2A}-(CH_2)_m-CO_2H$ , using one of the general methods that are described in the chemical literature, for example treating the acid with oxalyl chloride and reacting with  $R^L$  = hydroxy in a suitable solvent  
15 such as DMF.

Alternatively the above reactions may be carried out using a compound of formula (VIC):



20

(VIC)

- where Y and  $R^{1A}$  are as defined for formulae IIA and IIB and  $R^L$  is as defined for formulae (VIA) and (VIB)  
with the compound (VII) by the procedures (a), (b) or (c) set out above,  
and then  
25 treating the product with an acid,  
and where required or desired  
converting an  $R^{1A}$  or  $R^{2A}$  group to a  $R^1$  or  $R^2$  group, and/or  
converting one  $R^1$  or  $R^2$  group to another  $R^1$  or  $R^2$  group.

- 30 As mentioned previously, the acid treatment indicated above converts the epi-mutilin configuration of formula (VIC) to the usual mutilin nucleus of formula (VIA). Typically this conversion is carried out by treatment with conc. HCl or Lukas reagent (conc. HCl saturated with  $ZnCl_2$ ) in dioxane.

As in formulae (VIA) and (VIB),  $R^1A$  is typically the  $R^1$  group vinyl, and this may be converted to the alternative  $R^1$  group by hydrogenating the vinyl group to form an ethyl group. Also it may again be necessary to protect substituent groups in the compound (VII) prior to reaction, for example protecting N atoms with alkoxycarbonyl, for example *t*-butoxycarbonyl.

The compounds of formulae (VIA), (VIB) and (VIC) may be prepared by reacting the corresponding compounds of formula (IIA), (IIB) and (IIC) by conventional methodology to introduce acyl groups substituted by hydroxyl or amine or a leaving group.

Reference is directed to the preparation of the chloride and tosylate by K Riedl in *J. Antibiotics*, 1976, **29**, 132; and the tosylate and mesylate described by H Egger and H Reinshagen in *J. Antibiotics*, 1976, **29**, 915; starting from pleuromutilin or 19,20-dihydro-pleuromutilin ( $n=0$ ). Also compounds where  $R^L$  is chloro or bromo may be prepared by reacting  $Br(CH_2)_n(CH_2)COOCl$  or  $Cl(CH_2)_n(CH_2)COOCl$  with compounds IV and V above. It will be appreciated that when  $n=0$ , compounds where  $R^L$  is hydroxy are pleuromutilin and 19,20-dihydro-pleuromutilin. Compounds where  $R^L$  is  $NH_2$  may be prepared from the compound where  $R^L$  is a leaving group, for example treating a tosylate with sodium azide, followed by treatment with triphenyl phosphine and a base.

Compounds of formula (IA) wherein X is S(O) or  $SO_2$  may be obtained by preparing the corresponding compound in which X = S and treating it with an oxidising agent; for example, 3-chloroperoxybenzoic acid in chloroform, or catalytic osmium tetroxide plus *N*-methylmorpholine *N*-oxide in tetrahydrofuran and tertiary-butanol.

It will be appreciated that it is also possible to carry out the reaction of the compounds VIA/B/C with compound VII with the substituents reversed, i.e. with  $-CH_2(CH_2)_nXH$  as a 14-mutilin substituent and  $R^L$  on the  $R^{2A}-(CH_2)_m-$  residue. For example 22-deoxy-22-sulfanyl-pleuromutilin (US Patent 4130709) may be reacted with a compound of formula  $R^{2A}-(CH_2)_m-R^L$ , where  $R^L$  is a leaving group, such as 4-MeC<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>O, MeSO<sub>2</sub>O, CF<sub>3</sub>SO<sub>2</sub>O, or Cl, in the presence of an inorganic base, such as sodium methoxide, sodium ethoxide, or sodium hydride, in a solvent such as 2-propanol, ethanol, methanol, or tetrahydrofuran.

The compounds (III) and (VII) are commercially available or may be formed by conventional methodology from compounds that are commercially available compounds or described in the literature.

Where intermediates disclosed for the above processes are novel compounds, they also form part of this invention.

5 The compounds of the present invention may contain a chiral centre, and therefore the products of the above processes may comprise a mixture of diastereoisomers or a single diastereoisomer. A single diastereoisomer may be prepared by separating such a mixture of diastereoisomers which has been synthesised using a racemic starting material, or by synthesis using an optically pure starting material.

10 The products of the processes of this invention may be in crystalline or non-crystalline form, and, if crystalline, may optionally be hydrated or solvated. When some of the compounds of this invention are allowed to crystallise or are recrystallised from organic solvents, solvent of crystallisation may be present in the crystalline product. This invention includes within its scope such solvates. Similarly, some of the compounds of this  
15 invention may be crystallised or recrystallised from solvents containing water. In such cases water of hydration may be present in the crystalline product. This invention includes within its scope stoichiometric hydrates as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation.

20 The compounds obtained according to the processes of the invention are suitably worked up to a substantially pure form, for example at least 50% pure, suitable at least 60% pure, advantageously at least 75% pure, preferably at least 85% pure, more preferably at least 95% pure, especially at least 98% pure, all percentages being calculated as weight/weight. An impure or less pure form of a compound according to the invention may, for example,  
25 be used in the preparation of a more pure form of the same compound or of a related compound (for example a corresponding derivative) suitable for pharmaceutical use.

The present invention also includes pharmaceutically acceptable salts and derivatives of the compounds of the invention. Salt formation may be possible when one of the substituents  
30 carries an acidic or basic group. Salts may be prepared by salt exchange in conventional manner

Acid-addition salts may be pharmaceutically acceptable or non-pharmaceutically acceptable. In the latter case, such salts may be useful for isolation and purification of the  
35 compound of the invention, or intermediates thereto, and will subsequently be converted into a pharmaceutically acceptable salt or the free base. Pharmaceutically acceptable acid-addition salts include those described by Berge, Bighley, and Monkhouse, *J. Pharm. Sci.*, 1977, 66, 1-19. Suitable salts include the hydrochloride, maleate, and methanesulphonate; particularly the hydrochloride.



It will also be understood that where the compound of the invention contains a free carboxy moiety, it can form a zwitterion.

5 The compounds of the present invention and their pharmaceutically acceptable salts or derivatives have antimicrobial properties and are therefore of use in therapy, in particular for treating microbial infections in animals, especially mammals, including humans, in particular humans and domesticated animals (including farm animals). The compounds may be used for the treatment of infections caused by, for example, Gram-positive and  
10 Gram-negative bacteria and mycoplasmas, including, for example, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Enterococcus faecalis*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Haemophilus sp.*, *Neisseria sp.*, *Legionella sp.*, *Chlamydia sp.*, *Moraxella catarrhalis*, *Mycoplasma pneumoniae*, and *Mycoplasma gallisepticum*.

15 The present invention also provides a method of treating microbial infections in animals, especially in humans and in domesticated mammals, which comprises administering a compound of the invention or a pharmaceutically acceptable salt or derivative or solvate thereof, or a composition according to the invention, to a patient in need thereof.

20 The invention further provides the use of a compound of the invention or a pharmaceutically acceptable salt or derivative or solvate thereof in the preparation of a medicament for use in the treatment of microbial infections.

25 Compounds of the present invention may be used to treat skin and soft tissue infections and acne, by topical application. Accordingly, in a further aspect the present invention provides the use of a compound of the invention or a pharmaceutically acceptable salt or derivative or solvate thereof in the preparation of a medicament adapted for topical administration for use in the treatment of skin and soft tissue infections and also in the  
30 treatment of acne in humans.

Compounds of the present invention may be also used for the elimination or reduction of nasal carriage of pathogenic bacteria such as *S. aureus*, *H. influenzae*, *S. pneumonia* and *M. catarrhalis*, in particular colonisation of the nasopharynx by such organisms, by the  
35 administration of a compound of the present invention thereto. Accordingly, in a further aspect, the present invention provides for the use of a compound of the invention or a pharmaceutically acceptable salt or derivative or solvate thereof in the manufacture of a medicament adapted for administration to the nasal cavity, for reducing or eliminating the

nasal carriage of pathogenic organisms. Preferably, the medicament is adapted for focussed delivery to the nasopharynx, in particular the anterior nasopharynx.

Such reduction or elimination of nasal carriage is believed to be useful in prophylaxis of recurrent acute bacterial sinusitis or recurrent otitis media in humans, in particular in reducing the number of episodes experienced by a patient over a given period of time or reducing the time intervals between episodes. Accordingly, in a further aspect, the present invention provides for the use of a compound of the invention or a pharmaceutically acceptable salt or derivative or solvate thereof in the manufacture of a medicament adapted for administration to the nasal cavity, for prophylaxis of recurrent acute bacterial sinusitis or recurrent otitis media.

Compounds of the present invention are also useful in treating chronic sinusitis. Accordingly, in a further aspect, the present invention provides for the use of a compound of the invention or a pharmaceutically acceptable salt or derivative or solvate thereof in the manufacture of a medicament, for treating of chronic sinusitis.

The compounds according to the invention may suitably be administered to the patient at a daily dosage of from 1.0 to 50 mg/kg of body weight. For an adult human (of approximately 70 kg body weight), from 50 to 3000 mg, for example about 1500 mg, of a compound according to the invention may be administered daily. Suitably, the dosage for adult humans is from 5 to 20 mg/kg per day. Higher or lower dosages may, however, be used in accordance with normal clinical practice.

To lessen the risk of encouraging the development of resistant organisms during prophylaxis of recurrent otitis media or recurrent acute bacterial sinusitis, it is preferred to administer the drug on an intermittent, rather than a continual, basis. In a suitable intermittent treatment regimen for prophylaxis of recurrent otitis media or recurrent sinusitis, drug substance is administered on a daily basis, for a small number of days, for instance from 2 to 10, suitably 3 to 8, more suitably about 5 days, the administration then being repeated after an interval, for instance, on a monthly basis over a period of months, for instance up to six months. Less preferably, the drug substance may be administered on a continuing, daily basis, over a prolonged period, for instance several months. Suitably, for prophylaxis of recurrent otitis media or recurrent sinusitis, drug substance is administered once or twice a day. Suitably, drug substance is administered during the winter months when bacterial infections such as recurrent otitis media and recurrent sinusitis tend to be more prevalent. The drug substance may be administered at a dosage of from 0.05 to 1.00mg, typically about 0.1 to 0.2mg, in each nostril, once or twice a day.

More generally, the compounds and compositions according to the invention may be formulated for administration in any convenient way for use in human or veterinary medicine, by analogy with other antibiotics.

- 5 Accordingly, in a further aspect, the present invention provides a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt or derivative or solvate thereof together with a pharmaceutically acceptable carrier or excipient.
- 10 The compounds and compositions according to the invention may be formulated for administration by any route, for example oral, topical or parenteral. The compositions may, for example, be made up in the form of tablets, capsules, powders, granules, lozenges, creams, syrups, sprays or liquid preparations, for example solutions or suspensions, which may be formulated for oral use or in sterile form for parenteral administration by injection
- 15 or infusion.

Tablets and capsules for oral administration may be in unit dosage form, and may contain conventional excipients including, for example, binding agents, for example, syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar,

20 maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants, for example potato starch; and pharmaceutically acceptable wetting agents, for example sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice.

25 Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or another suitable vehicle before use. Such liquid preparations may contain conventional additives, including, for example, suspending agents, for

30 example sorbitol, methyl cellulose, glucose syrup, gelatin, hydroxyethyl cellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, oily esters (for example glycerine), propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl

35 p-hydroxybenzoate or sorbic acid; and, if desired, conventional flavouring and colour agents.

Compositions according to the invention intended for topical administration may, for example, be in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops.

nose drops, nasal sprays, impregnated dressings, and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, ethanol or  
5   oleyl alcohol for lotions and aqueous bases for sprays. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

10   Compositions according to the invention intended for topical administration, in addition to the above, may also contain a steroidal anti-inflammatory agent; for example, betamethasone.

15   Compositions according to the invention may be formulated as suppositories, which may contain conventional suppository bases, for example cocoa-butter or other glycerides.

20   Compositions according to the invention intended for parenteral administration may conveniently be in fluid unit dosage forms, which may be prepared utilizing the compound and a sterile vehicle, water being preferred. The compound, depending on the vehicle and concentration used, may be either suspended or dissolved in the vehicle. In preparing  
25   solutions, the compound may be dissolved in water for injection and filter-sterilised before being filled into a suitable vial or ampoule, which is then sealed. Advantageously, conventional additives including, for example, local anaesthetics, preservatives, and buffering agents can be dissolved in the vehicle. In order to enhance the stability of the solution, the composition may be frozen after being filled into the vial, and the water  
30   removed under vacuum; the resulting dry lyophilised powder may then be sealed in the vial and a accompanying vial of water for injection may be supplied to reconstitute the liquid prior to use. Parenteral suspensions may be prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilisation cannot be accomplished by filtration. The compound may instead be sterilised  
35   by exposure to ethylene oxide before being suspended in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in such suspensions in order to facilitate uniform distribution of the compound.

35   A compound or composition according to the invention is suitably administered to the patient in an antimicrobially effective amount.

A composition according to the invention may suitably contain from 0.001% by weight, preferably (for other than spray compositions) from 10 to 60% by weight, of a compound

according to the invention (based on the total weight of the composition), depending on the method of administration.

5 When the compositions according to the invention are presented in unit dosage form, for instance as a tablet, each unit dose may suitably comprise from 25 to 1000 mg, preferable from 50 to 500 mg, of a compound according to the invention.

10 Preferred compositions of the present invention include those adapted for intranasal administration, in particular, those that will reach into the nasopharynx. Such compositions are preferably adapted for focussed delivery to, and residence within, the nasopharynx. The term 'focussed delivery' is used to mean that the composition is delivered to the nasopharynx, rather than remaining within the nares. The term 'residence' within the nasopharynx is used to mean that the composition, once delivered to the nasopharynx, remains within the nasopharynx over a course of several hours, rather than  
15 being washed away more or less immediately. Preferred compositions include spray compositions and creams. Representative spray compositions include aqueous compositions, as well as oily compositions which contain amphiphilic agents so that the composition increases in viscosity when in contact with moisture. Creams may also be used, especially creams having a rheology that allows the cream to spread readily in the  
20 nasopharynx.

Preferred aqueous spray compositions include, in addition to water, further excipients including a tonicity modifier such as a salt, for instance sodium chloride; preservative, such as benzalkonium salt; a surfactant such as a non-ionic surfactant, for instance a  
25 polysorbate; and buffer, such as sodium dihydrogen phosphate; present in low levels, typically less than 1%. The pH of the composition may also be adjusted, for optimum stability of the drug substance during storage. For compounds of the present invention, a pH in the range 5 to 6, preferably about 5.3 to 5.8, typically about 5.5 is optimal

30 Representative oily spray and cream compositions are described in WO 98/14189 (SmithKline Beecham).

Suitably, the drug substance is present in compositions for nasal delivery in between 0.001 and 5%, preferably 0.005 and 3%, by weight of the composition. Suitable amounts include  
35 0.5% and 1% by weight of the composition (for oily compositions and creams) and from 0.01 to 0.2% (aqueous compositions).

Preferably, an aqueous spray composition is used. Such compositions are found to show similar retention in the target area (nasal cavity and nasopharynx) in gamma scintigraphy

studies and have superior release rates in synthetic membrane diffusion studies when compared to an oily composition as described in WO 98/14189. In addition, an aqueous base was found to be preferred to an oily base in sensory analysis studies.


- 5    Spray compositions according to the present invention may be delivered to the nasal cavity by spray devices well known in the art for nasal sprays, for instance an air lift pump. Preferred devices include those which are metered to provide a unit volume of composition, preferably about 100 $\mu$ l, and optionally adapted for nasal administration by addition of a modified nozzle.

10


The following Examples illustrate the present invention and particularly the preparative procedures outlined above, by reference to the preparation of specific compounds within the scope of the present invention.

### Note on naming of pleuromutilin analogues

In the Examples, compound (a), which in the IUPAC system has the systematic name (1*S*, 2*R*, 3*S*, 4*S*, 6*R*, 7*R*, 8*R*, 14*R*)-3,6-dihydroxy-2,4,7,14-tetramethyl-4-vinyl-tricyclo[5.4.3.0<sup>1,8</sup>]tetradecan-9-one, is referred to using the trivial name mutilin and with the numbering system described by H Berner, G Schulz, and H Schneider in *Tetrahedron*, 1981, 37, 915-919.




(a) IUPAC numbering



(a) Mutilin numbering

Likewise, compound (b), which has the systematic name (1*R*, 2*R*, 4*S*, 6*R*, 7*R*, 8*S*, 9*R*, 14*R*)-6-hydroxy-9-methoxy-2,4,7,14-tetramethyl-4-vinyl-tricyclo[5.4.3.0<sup>1,8</sup>]tetradecan-3-one, is named as (3*R*)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-*epi*-mutilin.



(b)

### Example 1 - Mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate

Quinuclidin-4-thiol hydrobromide (1.9 g, 0.009 mole) (W. Eckhardt and E.A. Grob, *Helvetica Chimica Acta* (1974), 57 (8)m 2339-2345) was added to a stirred solution of sodium ethoxide (1.72 g, 0.0253 mole) in ethanol (50 ml) under argon at room temperature. The mixture was stirred for 10 minutes before adding a solution of mutilin 14-toluenesulfonyloxyacetate (K. Ridel, *J. Antibiotics* (1976), 29m 132-139) (6.23 g, 0.0117 mole) in methyl ethyl ketone (20 ml). The mixture was stirred overnight at room temperature under argon, then concentrated *in vacuo*. The residue was partitioned between dichloromethane and water. The organic layer was washed with water, dried over magnesium sulphate, and concentrated *in vacuo*. The crude product was purified by chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution (19:1:0.1) to give the title compound as a solid 1.8 g (40%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia*

0.75 (3H, d, J 6.7 Hz), 0.88 (3H, d, J 7 Hz), 1.25 (3H, s), 1.46 (3H, s), 1.68 (6H, t, J 7.6 Hz), 2.93 (6H, t, J 7.6 Hz), 3.18 (2H, ABq), 3.35 (1H, m), 5.19 (1H, dd, J 17.5 and ), 5.33 (1H, dd), 6.45 (1H, dd, J 17.4 and 11 Hz). MS (EI) m/z 504 (M<sup>+</sup>)

5 **Example 2 - Mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate Hydrochloride**

Mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate (1.0 g) was dissolved in a minimum volume of acetone and a 1M solution of HCl in ether was added. The heterogeneous mixture was concentrated *in vacuo*. The residue was triturated with ether (20 ml) and 1M HCl/ether (5 ml) to give the title compound as a beige solid (0.94 g); <sup>1</sup>H NMR (D<sub>2</sub>O) *inter alia* 0.63

10 (3H, d, J 6 Hz), 0.86 (3H, d, J 6.8 Hz), 1.09 (3H, s), 1.36 (3H, s), 2.05 (6H, m), 3.40 (6H, m), 3.49 (1H, m), 5.10 (2H, m), 5.64 (1H, d, J 8.3 Hz), 6.29 (1H, dd, J 17.4 and 11 Hz). MS (EI) m/z 504 (M<sup>+</sup>)

**Example 3 - 19,20-Dihydromutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate**

15 A solution of mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate (0.314 g, 0.00063 mole) in ethanol (30 ml) was hydrogenated over 10% Pd-C paste (50% moisture content) at room temperature for 1 hour. The catalyst was filtered off and the filtrate concentrated *in vacuo*. The residue was dissolved in chloroform and washed with saturated aqueous sodium carbonate and dried over magnesium sulphate. The resulting solution was evaporated to dryness *in vacuo* to give the title compound (0.18 g) (57%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia*

20 0.71 (3H, 1d, J 6.6 Hz), 0.78 (3H, t, J 7.5 Hz), 0.94 (3H, d, J 7 Hz), 0.96 (3H, s), 1.43 (3H, s), 1.7 (6H, t, J 7 Hz), 2.40 (1H, m), 2.94 (6H, t, J 7.5 Hz), 3.19 (2H, s), 3.41 (1H, d, J 8.4 Hz), 5.60 (1H, d). MS (EI) m/z 506 (M<sup>+</sup>)

25 **Example 4 - Mutilin 14-(quinuclidin-3-yloxy)-acetate hydrochloride**

3-Quinuclidinol (0.635 g) in dry DMF (4 ml) was stirred under argon and treated with sodium hydride (0.21 g of a 60% dispersion in oil). After 1 hour the mixture was cooled to -15°C and a solution of mutilin 14-methanesulfonyloxyacetate (2.28 g, see H. Egger and H. Reinshagen, J. Antibiotics **29** (9), 915) in dry DMF (4 ml) was added dropwise. The

30 mixture was allowed to warm gradually to room temperature, left 1 hour and diluted with water (30 ml) and chloroform (30 ml). The layers were shaken and separated, the organic phase washed twice more with water, dried over MgSO<sub>4</sub> and evaporated. The residue was chromatographed on silica, eluting with dichloromethane/methanol/35% ammonia solution (19:1:0.1) to isolate a compound at R<sub>f</sub> approx. .45 on silica tlc, eluting with the same

35 solvent mixture. A solution of this compound in chloroform (5 ml) was treated with 1N HCl in ether (2 ml) and evaporated to provide the title compound as a buff foam (0.339 g);  $\nu_{\max}$  (CHCl<sub>3</sub>) 3562, 3435 (broad), 2447 (broad), 1735 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.71 (3H, d, J 6.7 Hz), 0.90 (3H, d, J 6.7 Hz), 3.1-3.6 (7H, m), 3.8-4.1 3H, m), 5.22 (1H, d,



17.5 Hz), 5.38 (1H, d, J 10.8 Hz), 5.81 (1H, d, J 8.3 Hz), 6.48 (1H, dd, J 14.7 and 11.0 Hz), 12.3 (1H, broad s, disappears on D<sub>2</sub>O exchange); MS (+ve ion electrospray) m/z 488 (MH<sup>+</sup>, 90%), 186 (100%).

#### 5 Example 5 - Mutilin 14-(quinuclidin-3-ylsulfanyl)-acetate

The preparation of quinuclidin-3-thiol was based on patent literature (J. Barriere, C. Cotret and J. Paris, E.P. 248703 [1987]). A solution of triphenylphosphine (12 g) in THF (85 ml) was ice-cooled under argon and treated dropwise with diisopropyl azodicarboxylate (9 ml). After 30 minutes a solution of 3-quinuclidinol (2.9 g) and thiolacetic acid (3.24 ml) in THF (170 ml) was added dropwise over 1 hour. The mixture was stirred overnight at room temperature, evaporated and the residue taken up in ether (250 ml). This solution was extracted with 1M hydrochloric acid (2 x 40 ml), the combined aqueous extracts washed with ether (100 ml) and evaporated to dryness. The residue was desiccated under vacuum over P<sub>2</sub>O<sub>5</sub> for 4 days to provide a pale yellow solid. A portion of this solid (0.443 g) was dissolved in ethanol (10 ml) and treated with sodium methoxide (0.216 g). After 1 hour, mutilin 14-methanesulfonyloxyacetate (0.912 g) was added, the mixture stirred a further 1 hour, diluted with chloroform (30 ml) and water (30 ml), shaken and separated. The organic layer was washed with water (30 ml), dried over MgSO<sub>4</sub> and evaporated. Chromatography of the residue on silica eluting with chloroform/methanol/35% ammonia solution (19:1:0.1) provided the title compound as a pale yellow foam, 0.62 g (62%);  $\nu_{\max}$  (CHCl<sub>3</sub>) 3563, 1730 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.74 (3H, d, J 6 Hz), 0.88 (3H, d, J 7 Hz), 5.1-5.4 (2H, m), 5.76 and 5.77 (1H, 2d, J 8.3 Hz), 6.49 (1H, dd, J 17 and 11 Hz); MS (+ve ion electrospray) m/z 504 (MH<sup>+</sup>, 100%), 202 (55%).

#### 25 Example 6 - Mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate

##### Step 1. Quinuclidin-4-thiol hydrochloride

Crude quinuclidin-4-thiol hydroiodide (Eckharat *et al.*, Helv. Chem. Acta, **57** (4), (1974) 2339-2345 (15.1 g, 0.057 mole) was dissolved in water (200 ml). Sodium carbonate (21.0 g, 0.2 mole) was added. The mixture was extracted with chloroform (200 ml x 7). The combined organic extract was dried over MgSO<sub>4</sub> and concentrated *in vacuo*. To the concentrate was added 1M hydrogen chloride in ether (100 ml). The mixture was evaporated to dryness *in vacuo* to yield the title compound as a white solid 7.135 g (71%); <sup>1</sup>H NMR (D<sub>2</sub>O) 2.18 (6H, t, J 8 Hz), 3.40 (6H, t, J 8 Hz), MS (EI) m/z 144 [(M-HCl)H]<sup>+</sup>, 100%).

##### 35 Step 2 Mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate

Quinuclidin-4-thiol hydrochloride (5 g) was stirred with ethanol (110 ml) under argon and solid sodium methoxide (3.15 g) added. After 30 minutes mutilin 14-methanesulfonyloxyacetate (12.7 g) was added, followed by ethanol (30 ml). After a further 30 minutes the mixture was diluted with chloroform (250 ml) and water (250 ml),

shaken and separated. The organic layer was washed with water (200 ml), dried over  $\text{MgSO}_4$  and evaporated. Chromatography of the residue on silica, eluting with chloroform/methanol/35% ammonia solution (19:1:0.1) provided the title compound as a pale coloured foam (12.24 g), identical by NMR with the product of Example 1.

5

**Example 7 - Mutilin 14-[N-(2,2-dimethylazabicyclo[4.3.0]non-4-ylmethyl)]-aminoacetate**

**Step 1 (+) Equatorial 4-cyano-2,2-dimethylazabicyclo[4.3.0]nonane**

To a mixture of (+) 2,2-dimethylazabicyclo[4.3.0] non-4-one (4.7 g, 0.028 mole), (F.D. King, J. Chem. Soc. Perkins. Trans 1, 447, 1986) and tosylmethylisocyanide (6.47 g, 0.033 mole) in dry dimethoxyethane (100 ml) at  $-10^\circ\text{C}$  was added ethanol (3.4 ml) followed by potassium-*tert*-butoxide (7.21 g, 0.064 mole). The mixture was stirred at  $-10^\circ\text{C}$  for 1 hour then warmed to  $50^\circ\text{C}$  for 2 hours. The mixture was allowed to cool and diethyl ether (500 ml) added. Filtration and concentration of the filtrate *in vacuo* gave an oil. Column chromatography on silica gel eluting with ethyl acetate gave the title compound as an oil 3.0 g (60%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 0.95 (3H, s), 1.21 (3H, s), 1.35-1.51 (2H, m), 1.61-1.91 (4H, m), 2.15-2.19 (1H, m), 2.28-2.39 (2H, m), 2.57-2.71 (1H, m), 2.89-2.98 (1H, m).

15

**Step 2 (+) Equatorial aminomethyl-2,2-dimethylazabicyclo[4.3.0]nonane**

(+) Equatorial 4-cyano-2,2-dimethylazabicyclo [ 4.3.0] nonane (1.0 g, 0.0056 mole) in tetrahydrofuran (50 ml) was treated with lithium aluminium hydride (1.07 g, 0.028 mole) and stirred at ambient temperature for 18 hours. Diethyl ether (50 ml) was then added followed by a mixture of water ( 4 ml) and 10% aqueous sodium hydroxide solution (1.5 ml). Filtration and concentration of the filtrate *in vacuo* gave the title compound 0.97 g (95%) as an oil;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 0.95 (3H, s), 1.20 (3H, s), 1.25-1.95 (9H, m), 2.25-2.40 (2H, m), 2.55 (2H, d, J 6 Hz), 2.89-2.97 (1H, m).

25

**Step 3 Mutilin 14-[N-(2,2-Dimethylazabicyclo[4.3.0]non-4-ylmethyl)]-aminoacetate**

(+) Equatorial aminomethyl-2,2-dimethylazabicyclo [4.3.0]nonane (0.1 g, 0.0006 mole) was treated with mutilin 14-toluenesulfonyloxyacetate (0.25 g, 0.0005 mole), (K Riedl, J Antibiotics 29 (2), 133, 1976) and N,N-diisopropylethylamine (0.1 ml, 0.0006 mole) in ethanol (20 ml) and heated under reflux for 6 hours. The mixture was then concentrated *in vacuo* and the residue partitioned between saturated sodium hydrogen carbonate solution and dichloromethane. The organics were separated and dried ( $\text{Na}_2\text{SO}_4$ ). Chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution (90:9:1) gave the title compound 0.08 g, (31%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 0.71 (3H, d, J 6.5 Hz) 0.90 (3H, d, J 6.5 Hz), 0.95 (3H, s), 1.25-2.55 (38H, m), 2.85-2.97 (1H, m), 3.19-3.39 (2H, m), 5.15 (1H, d, J 16.5 Hz), 5.31 (1H, d, J 11.1 Hz), 5.78 (1H, d, J 8.6 Hz), 6.50 (1H, dd, J 15.0 and 11.1 Hz). MS (+ve ion electrospray) m/z 543 ( $\text{MH}^+$ , 100%).

35

**Example 8 - Mutilin 14-(quinuclidin-4-ylcarbonylamino)-acetate**

**Step 1. Mutilin 14-azidoacetate**

To a stirred solution of mutilin 14-toluenesulfonyloxyacetate (5.33 g, 0.01 mole) in acetone (50 ml) was added a solution of sodium azide (0.7 g, 0.011 mole) in water (6.5 ml). A solid precipitated briefly then redissolved. The homogenous mixture was stirred for 2 hours at ambient temperature then heated to reflux for 3 hours. The mixture was concentrated *in vacuo* to low volume then diluted with chloroform. The resulting solution was washed three times with water then dried over magnesium sulfate. Concentration *in vacuo* gave a pale yellow foam which was purified by chromatography on silica gel. Elution with ethyl acetate/hexane mixtures provided the title compound as a white foam 3.3 g (82%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.73 (3H, d, J 6.8 Hz), 0.89 (3H, d, J 7.1 Hz), 1.23 (3H, s), 1.47 (3H, s), 3.37 (1H, dd, J 10.7 and 6.6 Hz), 3.77 (2H, s), 5.22 (1H, dd, J 17.4 and 1.3 Hz), 5.38 (1H, dd, J 11 and 1.3 Hz), 5.86 (1H, d, J 8.5 Hz), 6.49 (1H, dd, J 17.4 and 11 Hz).

**Step 2. Mutilin 14-(triphenylphosphinimino)-acetate**

Triphenylphosphine (0.275 g, 0.00105 mole) was added to a stirred solution of mutilin 14-azidoacetate (0.404 g, 0.001 mole) in dichloromethane maintained under an atmosphere of argon. The solution rapidly became homogenous and a gas was evolved. Stirring was continued for 17 hours; the mixture was then concentrated *in vacuo* to give the title compound as a white solid, obtained by filtration after trituration in petroleum ether 0.638 g (100 %); MS (+ve ion electrospray) m/z 638 (MH<sup>+</sup>, 100%)

**Step 3. Mutilin 14-aminoacetate**

Mutilin 14-(triphenylphosphinimino)-acetate (1 g, 0.00157 mole) was suspended in ethanol (25 ml) and potassium hydroxide (0.175 g, 0.00314 mole) was added. The mixture was stirred for 17 hours during which time it became homogeneous. 2M hydrochloric acid (1.7 ml) was then added, stirring continued for ten minutes and the mixture concentrated *in vacuo*. The residue was taken up in 2M hydrochloric acid and the solution washed three times with dichloromethane. The aqueous phase was then layered with dichloromethane and the pH adjusted to 11 by addition of solid potassium carbonate with vigorous stirring. The organic phase was then separated, the aqueous phase extracted with dichloromethane, the combined organic extract washed with brine, dried over magnesium sulfate and concentrated *in vacuo*. The title compound was obtained as a white foam 0.505 g (85 %); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.71 (3H, d, J 6.5 Hz), 0.89 (3H, d, J 6.9 Hz), 1.17 (3H, s), 1.45 (3H, s), 3.33 (3H, m), 5.21 (1H, d, J 17.4 Hz), 5.36 (1H, d, J 11 Hz), 5.78 (1H, d, J 8.4 Hz), 6.52 (1H, dd, J 17.4 and 11 Hz).

**Step 4. Quinuclidin-4-ylcarbonyl chloride hydrochloride**

Quinuclidine-4-carboxylic acid hydrochloride (0.192 g, 0.001 mole) was suspended in dichloromethane (5 ml) and dimethylformamide (1 drop) and oxalyl chloride (0.436 ml, 0.635 g, 0.005 mole) were added. The resulting suspension was heated to reflux under an atmosphere of argon for six hours. Following concentration of the suspension *in vacuo* the

residue was suspended in dichloromethane, concentrated *in vacuo* and finally dried *in vacuo* to give the title compound as a pale brown solid.

**Step 5. Mutilin 14-(quinuclidin-4-ylcarbonylamino)-acetate**

- Quinuclidin-4-ylcarbonyl chloride hydrochloride (0.001 mole theoretical, Step 4) was  
 5 suspended in dichloromethane (6 ml) and mutilin 14-aminoacetate (0.126 g, 0.00033 mole) was added. To the stirred suspension, under an atmosphere of argon, was added triethylamine (0.278 ml, 0.202 g, 0.002 mole) and stirring continued for 18 hours. Chloroform and water were added and the pH of the aqueous phase adjusted to 11 by addition of solid potassium carbonate. After shaking, the phases were separated, the  
 10 organic phase was washed once with saturated aqueous sodium hydrogen carbonate and once with brine, dried over magnesium sulfate and concentrated *in vacuo* to give the crude product as an off-white foam. Purification by chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution provided a pale yellow glass. The product was dissolved in 2M hydrochloric acid, the solution washed twice with dichloromethane,  
 15 then layered with dichloromethane. The pH of the aqueous phase was adjusted to 11 by addition of solid potassium carbonate. After shaking, the organic phase was separated, dried over magnesium sulfate and concentrated *in vacuo*. The residue was repeatedly dissolved in chloroform and concentrated *in vacuo*. Finally the residue was triturated with diethyl ether to give the title compound as a buff solid 0.0019 g (11 %); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  
 20 *inter alia* 0.71 (3H, d, J 6.9 Hz), 0.88 (3H, d, J 7 Hz), 1.18 (3H, s), 1.45 (3H, s), 2.96 (6H, m), 3.37 (1H, m), 3.93 (2H, d, J 4.9 Hz), 5.23 (1H, d, J 17.4 Hz), 5.36 (1H, d, J 11 Hz), 5.79 (1H, d, J 8.5 Hz), 6.02 (1H, m(br)), 6.47 (1H, dd, J 17.4 and 11 Hz); MS (+ve ion electrospray) m/z 515 (MH<sup>+</sup>, 100%).

**25 Example 9 - Mutilin 14-[(3R,4R)-Azabicyclo[2.2.1]hept-3-ylcarbonylamino]-acetate**

**Step 1. (3R,4R)-Azabicyclo[2.2.1]hept-3-ylcarbonyl chloride hydrochloride**

- 3R,4R-Azabicyclo[2.2.1]heptane-3-carboxylic acid hydrobromide (0.127 g, 0.0005 mole) was suspended in dichloromethane (2 ml) and dimethylformamide (1 drop) and oxalyl chloride (0.131 ml, 0.191 g, 0.0015 mole) were added. The mixture was stirred for 4 hours  
 30 under an atmosphere of argon. The resulting homogenous solution was concentrated *in vacuo*, the residue was dissolved in dichloromethane, concentrated *in vacuo* and finally dried *in vacuo* to give the title compound as an off-white solid.

**Step 2. Mutilin 14-[(3R,4R)-Azabicyclo[2.2.1]hept-3-ylcarbonylamino]-acetate**

- (3R,4R)-Azabicyclo[2.2.1]hept-3-ylcarbonyl chloride hydrochloride (0.0005 mole  
 35 theoretical, Step 1) was dissolved in dichloromethane (4 ml) and mutilin 14-aminoacetate (0.126 g, 0.00033 mole) was added. To the stirred solution under an atmosphere of argon was added triethylamine (0.134 ml, 0.101 g, 0.001 mole). The resulting solution was stirred for 17 hours. Chloroform and water were added and the pH of the aqueous phase adjusted to 11 by addition of solid potassium carbonate. After shaking, the phases were

separated, the organic phase was washed once with saturated aqueous sodium hydrogen carbonate and once with brine, dried over magnesium sulfate and concentrated *in vacuo* to give the crude product as an off-white foam. Purification by chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution provided the product as a white foam 0.142 g (86 %); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.72 (3H, d, J 6.9 Hz), 0.89 (3H, d, J 7 Hz), 1.18 (3H, s), 1.46 (3H, s), 3.37 (1H, m(br)), 3.96 (2H, d, J 5.1 Hz), 5.22 (1H, d, J 17.4 Hz), 5.36 (1H, d, J 11 Hz), 5.78 (1H, d, J 8.4 Hz), 5.96 (1H, m(br)), 6.47 (1H, dd, J 17.4 and 11 Hz); MS (+ve ion electrospray) m/z 501 (MH<sup>+</sup>, 40%).

#### 10 Example 10 - Mutilin 14-(1-methylpiperid-4-ylcarbonylamino)-acetate

##### Step 1. 1-Methylpiperid-4-ylcarbonyl chloride hydrochloride

1-Methylpiperidine-4-carboxylic acid hydrochloride (0.09 g, 0.0005 mole) was suspended in dichloromethane (5 ml) and dimethylformamide (1 drop) and oxalyl chloride (0.131 ml, 0.19 g, 0.0015 mole) were added. The mixture was stirred for 4 hours under an atmosphere of argon. The resulting homogenous solution was concentrated *in vacuo*, the residue was dissolved in dichloromethane, concentrated *in vacuo* and finally dried *in vacuo* to give the title compound as an off-white solid.

##### Step 2. Mutilin 14-(1-methylpiperid-4-ylcarbonylamino)-acetate

1-Methylpiperid-4-ylcarbonyl chloride hydrochloride (0.0005 mole theoretical, Step 1) was dissolved in dichloromethane (4 ml) and mutilin 14-aminoacetate (0.126 g, 0.00033 mole) was added. Triethylamine (0.139 ml, 0.101 g, 0.001 mole) was added to the stirred solution under an atmosphere of argon. After 2 hours chloroform and water were added and the pH of the aqueous phase adjusted to 11 by addition of solid potassium carbonate. After shaking, the phases were separated, the organic phase was washed once with saturated aqueous sodium hydrogen carbonate and once with brine, dried over magnesium sulfate and concentrated *in vacuo* to give the crude product as an off-white foam. Purification by chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution provided the product as a white foam 0.150 g (90 %); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.71 (3H, d, J 6.8 Hz), 0.89 (3H, d, J 7 Hz), 1.18 (3H, s), 1.45 (3H, s), 3.36 (1H, m), 3.94 (2H, d, J 5 Hz), 5.22 (1H, d, J 18.6 Hz), 5.35 (1H, d, J 12.2 Hz), 5.78 (1H, d, J 8.4 Hz), 5.99 (1H, m(br)), 6.47 (1H, dd, J 17.4 and 11 Hz); MS (+ve ion electrospray) m/z 503 (MH<sup>+</sup>, 25%).

#### Example 11 - Mutilin 14-[3-(1-methylpiperid-4-yl)]-propionate

##### 35 Step 1. 3-(1-Methylpiperid-4-yl)propionyl chloride

A suspension of 3-(1-methylpiperid-4-yl)propionic acid hydrochloride (WO 9620173 A1, Example 1)(0.33 g, 0.00159 mole) in dry dichloromethane (10 ml) was treated with dimethylformamide (1 drop) and oxalyl chloride (0.416 ml, 0.605 g, 0.00477 mole) under an atmosphere of argon. After stirring for 3½ hours the mixture was concentrated *in*

*vacuo*. The residue was dissolved in dry dichloromethane and concentrated *in vacuo* to give the title compound as a white solid.

**Step 2.**

A solution of 3-(1-methylpiperid-4-yl)propionyl chloride (0.00159 mole theoretical, Step 1) and (3R)-3-deoxy-11-deoxy-3-methoxy-11-oxo-4-*epi*-mutilin (H Berner, G. Schulz and H. Schneider, *Tetrahedron*, 1980, 36, 1807) in dry dimethylformamide was heated at 110°C under argon for 17 hours. The mixture was then concentrated *in vacuo* and the residue chromatographed on silica gel eluting with dichloromethane/methanol/35% ammonia solution mixtures. The title compound was obtained as a pale yellow oil 0.284 g (49%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.79 (3H, d, J 6.9 Hz), 0.99 (3H, d, J 6.4 Hz), 1.18 (3H, s), 1.24 (3H, s), 2.37 (3H, s), 2.97 (3H, m), 3.23 (3H, s), 3.48 (1H, m), 5.01 (1H, d, J 17.6 Hz), 5.30 (1H, d, J 10.7 Hz), 5.74 (1H, d, J 10 Hz), 6.67 (1H, dd, J 17.5 and 10.6 Hz); MS (+ve ion electrospray) m/z 488 (MH<sup>+</sup>, 100%).

**Step 3. Mutilin 14-[3-(1-methylpiperid-4-yl)]-propionate**

A solution of (3R)-3-deoxy-11-deoxy-3-methoxy-11-oxo-4-*epi*-mutilin (0.355 g, 0.000728 mole) in dioxan (3 ml) was treated with concentrated hydrochloric acid (3 ml). After 4 hours the mixture was diluted with water, layered with dichloromethane and the vigorously stirred mixture adjusted to pH 11 by addition of solid potassium carbonate. The phases were then separated and the aqueous phase extracted with dichloromethane. The combined organic extract was washed with saturated aqueous sodium hydrogen carbonate and brine, dried over magnesium sulfate and concentrated *in vacuo*. The residue was chromatographed on silica gel eluting with dichloromethane/methanol/35% ammonia solution (90:9:1) to give the title compound as a white foam 0.284 g (82%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.70 (3H, d, J 6.6 Hz), 0.88 (3H, d, J 7 Hz), 1.17 (3H, s), 1.45 (3H, s), 2.44 (3H, s), 3.05 (2H, m), 3.36 (1H, dd, J 11.4 and 7.4 Hz), 5.20 (1H, d, J 17.5 Hz), 5.36 (1H, d, J 11 Hz), 5.74 (1H, d, J 8.4 Hz), 6.51 (1H, dd, J 17.5 and 11 Hz); MS (+ve ion electrospray) m/z 474 (MH<sup>+</sup>, 100%).

**Example 12 - Mutilin 14-(quinuclid-4-ylmethylsulfanyl)-acetate**

An ice cooled solution of triphenylphosphine (1.19 g, 0.0042 mole) in dry tetrahydrofuran was treated dropwise with diisopropyl azodicarboxylate (0.85 g, 0.0042 mole). After 30 minutes a solution of quinuclid-4-ylmethanol (0.565 g, 0.004 mole) and thiolacetic acid (0.315 ml, 0.0042 mole) in dry tetrahydrofuran (20 ml) added dropwise over a period of 10 minutes. The mixture was left at 5° for 72 hours then concentrated *in vacuo* and the residue dissolved in ether (200 ml). The resulting solution was extracted with 1M hydrochloric acid (3 x 50 ml). The combined extract was concentrated *in vacuo* and dried *in vacuo* to give a gummy residue 0.65 g. The residue was dissolved in ethanol (30 ml) and treated under argon with potassium tert butoxide (0.785 g, 0.007 mole) for 30 minutes. Mutilin 14-methanesulfonyloxyacetate (1.38 g, 0.003 mole) was then added to the

ethanolic solution and the mixture stirred overnight under argon. The insoluble byproducts were filtered off and the filtrate evaporated to dryness. The residue was partitioned between chloroform and water. The organic layer washed with brine and dried over magnesium sulfate and evaporated to dryness. Chromatography of the residue on silica gel eluting with chloroform/methanol/35% ammonia solution (19:1:0.1) provided the title compound as a white foam, 0.48 g (31%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.74 (3H, d, J 6.6 Hz), 0.88 (3H, d, J 7 Hz), 1.76 (3H, s), 1.44 (6H, t, J 7.7 Hz), 2.47 (2H, s), 2.87 (6H, t, J 7.5 Hz), 3.09 (2H, s), 3.36 (1H, m), 5.1-5.4 (2H, m), 5.75 (1H, d, J 8.3 Hz), 6.48 (1H, m); MS (+ve ion electrospray) m/z 518 (MH<sup>+</sup>, 100%).

**Example 13 - 19,20-Dihydromutilin 14-(quinuclidin-4-ylsulfonyl)acetate**

19,20-Dihydromutilin 14-(quinuclidin-4-ylthio)acetate (0.05 g, 0.0001 mole) in dry tetrahydrofuran (2 ml) and tert-butanol (0.2 ml) was treated with N-methylmorpholine oxide (0.036 g, 0.003 mole) and a catalytic amount of osmium tetroxide under argon for a period of 4.5 hours. The mixture was extracted with ethyl acetate. The organic solution was filtered to remove inorganic residues. The filtrate was concentrated *in vacuo*.

Chromatography of the residue on silica gel eluting with chloroform/methanol/35% ammonia solution provided the title compound as a foam, 0.043 g (80%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.70 (3H, d, J 7 Hz), 0.82 (3H, t, J 7 Hz), 0.93 (3H, d, J 7 Hz), 0.96 (3H, s), 1.92 (6H, t, J 7.5 Hz), 3.01 (6H, t, J 7.5 Hz), 3.41 (1H, m), 3.73 (1H, d, J 13.3 Hz), 3.87 (1H, d, J 13.3 Hz), 5.68 (1H, d, J 8 Hz); MS (+ve ion electrospray) m/z 538 (MH<sup>+</sup>, 60%).

**Example 14 - 19,20-Dihydromutilin 14-(quinuclidin-4-ylsulfoxy)-acetate**

A cooled solution of 19,20-dihydromutilin 14-(quinuclidin-4-ylthio)-acetate (0.152 g, 0.0003 mole) and glacial acetic acid (0.06 g, 0.001 mole) in chloroform (5 ml) was treated with 80% 3-chloroperoxybenzoic acid (0.069 g, 0.0032 mole) at 0°C, allowed to warm to room temperature and stirred for 72 hours. The solvents were removed *in vacuo*.

Chromatography of the residue on silica gel eluting with chloroform/methanol/35% ammonia solution (20:1:0.1) provided the title compound as a white solid, 0.064 g (41%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.72-0.95 (12H, m), 1.45 (6H, t, J 8.5 Hz), 1.74 (6H, t, J 8 Hz), 2.13-2.25 (3H, m), 2.39 (1H, m), 3.02 (6H, t, J 7.6 Hz), 3.35-3.42 (3H, m), 5.71 (1H, d, J 8.4 Hz); MS (+ve ion electrospray) m/z 522 (MH<sup>+</sup>, 100%).

**Example 15 - Mutilin-14-(1-methylpiperid-4-ylsulfanyl)-acetate**

A solution of triphenylphosphine (5.51 g, 0.021 mole) in dry tetrahydrofuran (100 ml) was ice-cooled under argon and treated with diisopropyl azodicarboxylate (4.25 g, 0.021 mole). After 30 minutes a solution of 4-hydroxy-1-methylpiperidine (2.3 g, 0.02 mole) and thiolacetic acid (1.54 g, 0.02 mole) in dry tetrahydrofuran (50 ml) was added over a period

of 30 minutes. The mixture was stirred overnight at room temperature, evaporated *in vacuo* and the residue taken up in ether (200 ml). The ethereal solution was extracted with 1M hydrochloric acid (50 ml x 4). The combined aqueous extract was washed with ether, evaporated to dryness and dried *in vacuo* to give a yellow gum (2.4 g). A portion of this gum (0.517 g) was dissolved in ethanol and treated with potassium tert-butoxide (0.785 g) under argon for 30 minutes. Mutilin-14-methanesulfonyloxyacetate (0.92 g, 0.002 mole) was added and the mixture stirred overnight, then concentrated *in vacuo*. The residue was partitioned between chloroform and water. The organic layer was washed with brine, dried over magnesium sulfate and concentrated *in vacuo*. Chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution provided the title compound as a foam, 0.557 g (57%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.73 (3H, d, J 6.5 Hz), 0.87 (3H, d, J 7 Hz), 1.30 (3H, s), 1.67 (3H, s), 2.25 (3H, s), 3.16 (2H, s), 3.36 (1H, m), 5.28 (2H, m), 5.77 (1H, d, J 8.5 Hz), 6.47 (1H, m); MS (+ve ion electrospray) m/z 492 (MH<sup>+</sup>, 100%).

- Example 16 - Mutilin 14-((3RS,4SR)-1-aza-bicyclo[2.2.1]hept-3-yl-sulfanyl)-acetate**  
The title compound was prepared in 32% overall yield from *endo*-3-hydroxy-azabicyclo[2.2.1]heptane (S.M. Jenkins *et al*, *J. Med. Chem.*; 1992, 35, 2392-2406) using the procedure described in Example 5. The title compound was isolated as a colourless solid, in the <sup>1</sup>H NMR spectrum the 8 line multiplets at δ 3.05-3.40 and 6.43-6.56 indicate a 1:1 mixture of diastereoisomers; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.74 (3H, d, J 6.4 Hz), 0.88 (3H, d, J 7.0 Hz), 3.05-3.40 (2H, m), 5.21 (1H, d, J 17.5 Hz), 5.35 (1H, d, J 11.0 Hz), 5.75-5.80 (1H, m), 6.43-6.56 (1H, m); MS (+ve electrospray) m/z 490 (MH<sup>+</sup>).

- Example 17 - Mutilin 14-((3RS,4SR)-1-aza-bicyclo[2.2.1]hept-3-yl-sulfanyl)-acetate hydrochloride**  
The title compound was prepared from mutilin 14-((3RS,4SR)-1-azabicyclo[2.2.1]hept-3-yl-sulfanyl)-acetate using the procedure described in Example 2. The title compound was isolated as a colourless solid, 1:1 mixture of diastereoisomers; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) *inter alia* 0.65 (3H, d, J 6.4 Hz), 0.84 (3H, d, J 6.8 Hz), 1.09 (3H, s), 1.39 (1H, s), 4.61 (1H, d, J 5.2 Hz, exchanged with D<sub>2</sub>O), 5.05-5.12 (2H, m), 5.60 (1H, d, J 7.9 Hz), 6.14 (1H, dd, J 18 and 10.7 Hz), 10.4-10.6 (1H, br, exchanged with D<sub>2</sub>O); MS (+ve electrospray) m/z 490 (MH<sup>+</sup> of free base).

- Example 18 - Mutilin 14-(quinuclidin-3-ylidene)-acetate hydrochloride (both geometric isomers)**

**Step 1. Methyl quinuclidin-3-ylidene acetate hydrochloride**

A suspension of quinuclidin-3-one hydrochloride (3.23 g) in DMF (20 ml) was treated with sodium methoxide (1.08 g) and stirred vigorously for 30 minutes. A solution of trimethyl phosphonoacetate (4.05 ml) and sodium methoxide (1.35 g) in DMF (20 ml) was added



dropwise over 15 minutes and stirred a further 2½ hours. The DMF was evaporated and the residue treated with dry ether (100 ml), triturated and filtered. The filtrate was treated with 1N HCl in ether (30 ml), the resulting solid triturated and the ether decanted. Ether (200 ml) was added, the suspension stirred vigorously for 30 minutes, the solid filtered off and heated at 60°C under vacuum for 2 days. The resulting methylquinuclidin-3-ylideneacetate hydrochloride (3.93 g) was a ca. 1:1 mixture of geometric isomers; <sup>1</sup>H NMR (D<sub>2</sub>O) *inter alia* 5.84 (broad s) and 5.94 (t, J 2.5 Hz) (vinyl protons of the two geom. isomers).

**Step 2. Quinuclidin-3-ylideneacetic acid hydrochloride**

Methyl quinuclidin-3-ylideneacetate hydrochloride (1g) was heated in concentrated hydrochloric acid (10 ml) at 60°C for 18 hrs and the solution evaporated to dryness. The residue was kept under vacuum over P<sub>2</sub>O<sub>5</sub> for 3 days to give quinuclidin-3-ylidene acetic acid hydrochloride, 0.91 g (97%) as a white solid; <sup>1</sup>H NMR (D<sub>2</sub>O) *inter alia* 5.77 (broad s) and 5.86 (broad s) (ca. 1:1, vinyl protons of the two geom. isomers).

**Step 3. (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-(quinuclidin-3-ylidene)-acetate**

Quinuclidin-3-ylideneacetic acid hydrochloride (0.204 g) was suspended in chloroform (5 ml), stirred under argon and treated with 1 drop DMF and oxalyl chloride (0.87 ml). After 2 hours the solvent was evaporated, toluene (10 ml) added to the residue and evaporated.

The residue was taken up in DMF (2 ml), treated with (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin (0.334 g, prepared according to H. Berner, G. Schulz and H. Schneider, *Tetrahedron* (1980) 36 1807) and heated at 100°C under argon for 3 hours. After leaving at room temperature overnight, the mixture was diluted with chloroform (20 ml), washed with aqueous NaHCO<sub>3</sub> and water, dried and evaporated to dryness.

Chromatography on silica, eluting with chloroform/methanol/35% ammonia solution (19:1:0.1) separated the 2 geometric isomers of the title compound.

Less polar isomer, 0.1 g (20%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 3.23 (3H, s), 3.4-3.6 (1H, m), 3.96 (2H, ABq, J 20 Hz), 5.02 (1H, d, J 17.5 Hz), 5.34 (1H, d, J 10.5 Hz), 5.64 (1H, t, J 2.5 Hz), 5.81 (1H, d, J 10 Hz), 6.74 (1H, dd, J 17.5 and 10.5 Hz); MS (+ve ion electrospray) m/z 484 (MH<sup>+</sup>, 100%).

More polar isomer, 0.234 g (48%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 3.12 (2H, s), 3.23 (3H, s), 3.4-3.5 (1H, m), 5.01 (1H, d, J 17.5 Hz), 5.30 (1H, d, J 10.5 Hz), 5.78 (1H, d, J 10 Hz), 6.37 (1H, d, J 0.95 Hz), 6.65 (1H, dd, J 17.5 and 10.5 Hz); MS (+ve ion electrospray) m/z 484 (MH<sup>+</sup>, 100%).

**Step 4. Mutilin 14-(quinuclidin-3-ylidene)-acetate hydrochloride**

The less polar geometric isomer of (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-(quinuclidin-3-ylidene)-acetate (0.1 g) was dissolved in dioxan (3 ml), briefly cooled in ice-water and treated with conc. HCl (2 ml). After 5 hours at room temperature CHCl<sub>3</sub> (10 ml) and water (20 ml) were added, followed by solid NaHCO<sub>3</sub>

- until basic. The layers were separated, the aqueous re-extracted with  $\text{CHCl}_3$  and the combined organic dried and evaporated. The residue was chromatographed, eluting with chloroform/methanol/35% ammonia solution 97:3:0.3 and the product in chloroform solution treated with 1M HCl in ether (1 ml). Evaporation gave the less polar geometric isomer of the title compound as a white foam. 0.105 g;  $^1\text{H}$  NMR ( $\text{CD}_3\text{SOCD}_3$ ) *inter alia* 2.85 (1H, s), 4.38 (2H, ABq, J 19 Hz), 4.59 (1H, d, J 6 Hz, disappears on  $\text{D}_2\text{O}$  exchange), 5.0-5.2 (2H, m), 5.64 (1H, d, J 8 Hz), 5.92 (1H, s), 6.27 (1H, dd, J 17.5 and 11 Hz), 10.7 (1H, broad s, disappears on  $\text{D}_2\text{O}$  exchange); MS (+ve ion electrospray) m/z 470 ( $\text{MH}^+$  - HCl, 100%).
- 10 In the same manner, the more polar geometric isomer of (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-(quinuclidin-3-ylidene)-acetate (0.116 g) was converted into the more polar geometric isomer of the title compound (0.096 g) as an off-white foam;  $^1\text{H}$  NMR ( $\text{CD}_3\text{SOCD}_3$ ) *inter alia* 4.62 (1H, d, J 6 Hz, disappears on  $\text{D}_2\text{O}$  exchange), 5.0-5.2 (2H, m), 5.65 (1H, d, J 8 Hz), 6.18 (1H, dd, J 17.5 and 11 Hz), 6.64 (1H, s), 11.42 (1H, broad s, disappears on  $\text{D}_2\text{O}$  exchange); MS (+ve ion electrospray) m/z 470 ( $\text{MH}^+$  - HCl, 100%).

#### Example 19 - Mutilin 14-[(+)-quinuclidin-3-yl]-acetate hydrochloride

##### Step 1. (+)-Quinuclidine-3-acetic acid hydrochloride

- 20 A mixture of methyl quinuclidin-3-ylideneacetate hydrochloride (Example 18, Step 1) (2g), ethanol (50 ml), 2M hydrochloric acid (5 ml) and 10% Pd/C (1 g) was stirred for 24 hours under  $\text{H}_2$  at atmospheric pressure, filtered through celite and evaporated to dryness. The residue was dissolved in concentrated hydrochloric acid (10 ml), heated at  $60^\circ\text{C}$  for 18 hours, treated with a further 10 ml of concentrated hydrochloric acid, heated at  $80^\circ\text{C}$  for 6 hours and evaporated to dryness. The residue was kept under vacuum over  $\text{P}_2\text{O}_5$  for 3 days to give the title compound as a white solid (1.8 g); MS (+ve ion electrospray) m/z 170 ( $\text{MH}^+$ , 100%).

##### Step 2. (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-[(+)-quinuclidin-3-yl]-acetate

- 30 (+)-Quinuclidine-3-acetic acid hydrochloride was converted into acid chloride and reacted with (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin in the manner of Example 18, Step 3. Chromatography of the product gave the title compound as a white foam (68%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 3.23 (3H, s), 3.3-3.5 (1H, m), 5.01 (1H, d, J 17.5 Hz), 5.32 (1H, d, J 10.5 Hz), 5.75 (1H, d, J 9.8 Hz), 6.68 and 6.69 (1H, 2 dd, J 17.5 and 10.5 Hz); MS (+ve ion electrospray) m/z 486 ( $\text{MH}^+$ , 100%).

##### Step 3. Mutilin 14-[(+)-quinuclidin-3-yl]-acetate hydrochloride

Rearrangement of (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-[(+)-quinuclidin-3-yl]acetate in the manner of Example 18. Step 4 gave the title compound as a

white foam (95%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 5.1-5.4 (2H, m), 5.74 (1H, d, J 8.3 Hz), 6.43 and 6.47 (1H, 2 dd, J 17.5 and 10.5 Hz); MS (+ve ion electrospray) m/z 472 ( $\text{MH}^+$ , 100%).

5 **Example 20 - Mutilin 14-[(+)-quinuclidin-3-ylacetoxyl]-acetate hydrochloride**

(+)-Quinuclidine-3-acetic acid hydrochloride (0.206 g) was suspended in chloroform (5 ml) under argon, treated with DMF (1 drop) and oxalyl chloride (0.87 ml) and stirred 1 hour. The solution was evaporated, toluene was added and evaporated and the residue taken up in DMF (2 ml). Pleuromutilin (0.378 g) was added and the mixture stirred under argon for 18  
10 hours, then heated at 110°C for 30 minutes. It was diluted with chloroform (10 ml), washed with aqueous  $\text{NaHCO}_3$  (twice) and water, dried and evaporated. The residue was chromatographed, eluting with chloroform/methanol/35% ammonia solution (9:1:0.1). A chloroform solution of the material obtained was treated with 1M HCl in ether (2 ml) and evaporated. Trituration under ether and filtration gave the title compound as an off-white  
15 solid, 0.22 g (42%);  $^1\text{H}$  NMR ( $\text{CD}_3\text{SOCD}_3$ ) *inter alia* 4.5-4.7 (3H, m, reduces to 2H, m on  $\text{D}_2\text{O}$  exchange); 5.0-5.2 (2H, m), 5.59 (1H, d, J 8 Hz), 6.10 (1H, dd, J 17.5 and 10.5 Hz), 10.06 (1H, broad s, disappears on  $\text{D}_2\text{O}$  exchange); MS (+ve ion electrospray) m/z 530 ( $\text{MH}^+$ , 100%).

20 **Example 21 - Mutilin 14-(quinuclidin-3-ylmethylsulfanyl)-acetate**

**Step 1. Mixture of (+)-quinuclidin-3-ylmethylsulfanylacetate hydrochloride and (+)-quinuclidin-3-ylmethanethiol hydrochloride**

(+)-Quinuclidine-3-methanol (L.I. Mastafonova, L.N. Yakhontov, M.V. Rubtsov, Khim. Geterotsikl. Soedin., Akad. Nauk Latv. SSR. 1965(6), 858-863) was converted into the title  
25 mixture using the procedure of Example 5. MS (+ve ion electrospray) m/z 200 ( $\text{MH}^+$  for thioacetate, 100%), 158 ( $\text{MH}^+$  for thiol, 40%).

**Step 2. Mutilin 14-(+)-quinuclidin-3-ylmethylsulfanyl)-acetate**

The mixture from Step 1 was reacted with mutilin 14-methanesulfonyloxyacetate in the manner described in Example 5 to provide the title compound as an off-white foam (28%);  
30  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.75 (3H, d, J 6.7 Hz), 0.89 (3H, d, J 7.0 Hz), 3.12 (2H, s), 3.37 (1H, broad, becomes d, J 6.3 Hz on  $\text{D}_2\text{O}$  exchange), 5.21 (1H, d, J 17.5 Hz), 5.36 (1H, d, J 11 Hz), 5.75 (1H, d, J 8.4 Hz), 6.51 (1H, dd, J 17.5 and 11 Hz); MS (positive ion electrospray) m/z 518 ( $\text{MH}^+$ , 100%).

35 **Example 22 - 1,2-Didehydromutilin 14-(quinuclidin-4-ylsulfanyl)-acetate**

**Step 1. 1,2-Didehydromutilin-14-methanesulfonyloxyacetate**

1,2-Didehydropleuromutilin (0.2g, 0.00053 mole) (G. Schulz and H. Berner, *Tetrahedron*, (1984) 40, 905-17) was converted to 1,2-didehydromutilin-14-methanesulfonyloxyacetate by the method previously described for pleuromutilin (H. Egger and H. Reinshagen. *J.*

*Antibiotics* (1976), 29, 915-22) providing the product as a yellow foam, (100%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.80 (3H, d, J 6.7 Hz), 1.10 (3H, d, J 7.0 Hz), 1.16 (3H, s), 1.54 (3H, s), 3.21 (3H, s), 4.67 (2H, s), 5.22 (1H, dd, J 17.4 and 1.3 Hz), 5.38 (1H, dd, J 11 and 1.2 Hz), 5.81 (1H, d, J 8.9 Hz), 6.05 (1H, d, J 6.1 Hz), 6.44 (1H, dd, 17.3 and 11 Hz), 7.74 (1H, d, J 6.1 Hz).

**Step 2. 1,2-Didehydromutilin 14-(quinuclidin-4-ylsulfanyl)-acetate**

A solution of 1,2-didehydromutilin 14-methanesulfonyloxyacetate (0.00053 moles) in ethanol was treated with quinuclidin-4-thiol hydrochloride (0.105 g, 0.000583 mole). After 15 minutes sodium methoxide (0.057 g, 0.00106 mole) was added to the stirred solution. After 1 hour the mixture was concentrated *in vacuo* to a slurry. Chloroform and water were then added. The pH of the aqueous phase was adjusted to 11-12 by addition of solid potassium carbonate. The phases were separated and the aqueous re-extracted with chloroform. The combined organic extract was dried over magnesium sulfate and concentrated *in vacuo*. Purification by chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution provided the product as an off-white foam 0.19 g (72%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.80 (3H, d, J 6.4 Hz), 1.08 (3H, d, J 7 Hz), 1.15 (3H, s), 1.55 (3H, s), 3.20 (2H, ABq), 5.20 (1H, dd, J 17.4 and 1.4 Hz), 5.35 (1H, dd, J 11 and 1.4 Hz), 5.74 (1H, d, J 8.7 Hz), 6.04 (1H, d, J 6.1 Hz), 6.47 (1H, dd, J 17.3 and 11 Hz), 7.74 (1H, d, J 6.1 Hz); MS (-ve ion electrospray) m/z 500 ([M-H]<sup>-</sup>, 50%).

**Example 23 - 2α-Hydroxymutilin 14-(quinuclidin-4-ylsulfanyl)-acetate**

**Step 1. 2-Diazomutilin 14-methanesulfonyloxyacetate**

2-Diazopleuromutilin (0.809 g, 0.002 mole) (G. Schulz and H. Berner, *Tetrahedron* (1984), 40, 905-17) was converted to 2-diazomutilin-14-methanesulfonyloxyacetate by the method described for pleuromutilin (H. Egger and H. Reinshagen, *J. Antibiotics* (1976), 29, 915-22) providing the product as a bright yellow gum (100%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.75 (3H, d, J 6.9 Hz), 0.93 (3H, d, J 6.9 Hz), 1.18 (3H, s), 1.50 (3H, s), 3.20 (3H, s), 4.65 (2H, s), 5.24 (1H, d, J 17.5 Hz), 5.37 (1H, d, J 11 Hz), 5.84 (1H, d, J 8.5 Hz), 6.43 (1H, dd, J 17.4 and 11 Hz).

**Step 2. 2α-Dichloroacetoxymutilin-14-methanesulfonyloxyacetate**

2-Diazomutilin-14-methanesulfonyloxyacetate (0.002 moles theoretical) from Step 1 in dichloromethane (20 ml) was cooled in an ice bath under an atmosphere of argon. To the stirred solution was added dichloroacetic acid (0.309 g, 0.0024 mole), dropwise over 2 minutes. Stirring was continued for 2.5 hours. The mixture was diluted with dichloromethane and washed twice with saturated aqueous sodium hydrogen carbonate and once with brine. After drying over magnesium sulfate concentration *in vacuo* gave the product as a pale yellow foam (100%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.76 (3H, d, J 6 Hz), 0.93 (3H, d, J 7 Hz), 1.12 (3H, s), 1.49 (3H, s), 3.20 (3H, s), 4.66 (2H, s), 5.05 (1H, t, J 9

Hz), 5.25 (1H, d, J 17.3 Hz), 5.38 (1H, d, J 11 Hz), 5.83 (1H, d, J 8.5 Hz), 5.97 (1H, s), 6.43 (1H, dd, J 17.4 and 11 Hz).

**Step 3. 2 $\alpha$ -Hydroxymutilin 14-(quinuclidin-4-ylsulfanyl)-acetate**

2-Dichloroacetoxymutilin-14-methanesulfonyloxyacetate (0.001 mole theoretical) from  
5 Step 2 in ethanol (2 ml) was added to a pre-mixed solution of quinuclidine-4-thiol  
hydrochloride (0.27 g, 0.0015 mole) and sodium methoxide (0.162 g, 0.003 mole) in  
ethanol (8 ml). After stirring for 1 hour the mixture was diluted with chloroform, washed  
twice with saturated sodium hydrogen carbonate and once with brine, then dried over  
magnesium sulfate. Concentration *in vacuo* was followed by silica gel chromatography,  
10 eluting with chloroform/methanol/35% ammonia solution. The product was obtained as a  
white foam 0.2 g (38% overall, 3 steps); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.75 (3H, d, J 6.5  
Hz), 0.92 (3H, d, J 7 Hz), 1.17 (3H, s), 1.48 (3H, s), 3.19 (2H, ABq), 3.99 (1H, t, J 8.7 Hz),  
5.20 (1H, d, J 17.3 Hz), 5.33 (1H, d, J 11 Hz), 5.75 (1H, d, J 8.4 Hz), 6.45 (1H, dd, J 17.3  
and 11 Hz); MS (+ve ion electrospray) m/z 520 (MH<sup>+</sup>, 100%).

15

**Example 24 - Mutilin 14-(quinuclidin-4-yl)-acetate**

**Step 1. Quinuclidin-4-ylmethanol**

Quinuclidine-4-carboxylic acid hydrochloride (6.0 g, 0.031 mmoles) in tetrahydrofuran  
(300 ml) was treated with lithium aluminium hydride (5.0 g, 0.137 mmoles) at ambient  
20 temperature for 18 hours. Water (20 ml) and 10% aqueous sodium hydroxide (7.5 ml) was  
added carefully and the mixture filtered, washing with diethyl ether. The combined  
filtrates were evaporated to dryness to give the title compound as a white solid 4.04 g,  
(91%); MS (+ve ion electrospray) m/z 142 (MH<sup>+</sup>, 100%)

**Step 2. Quinuclidin-4-ylacetonitrile**

25 Quinuclidin-4-ylmethanol (2.19 g, 0.015 moles) was converted to the corresponding  
mesylate by treatment with triethylamine/methanesulphonyl chloride in chloroform.  
Washing the organics with saturated potassium carbonate, drying over sodium sulphate and  
evaporation to dryness gave the mesylate 3.24 g (95%). The mesylate was dissolved in dry  
dimethyl formamide (50 ml) and treated with sodium cyanide (2.26 g, 0.046 moles) and  
30 heated to 130°C for 18 hours. The mixture was evaporated to dryness and the residue  
partitioned between saturated potassium carbonate and chloroform. The organics were  
dried (Na<sub>2</sub>SO<sub>4</sub>) and chromatographed on silica gel eluting with 0-10%  
methanol/chloroform. This gave the title compound 1.1 g (50%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.45  
(6H, t, J 9 Hz), 2.12 (2H, s), 2.85 (6H, t, J 9 Hz); MS (+ve ion electrospray) m/z 151  
35 (MH<sup>+</sup>, 100%).

**Step 3. Ethyl quinuclidin-4-ylacetate**

Hydrogen chloride gas was bubbled through a solution of quinuclidin-4-ylacetonitrile (1.1  
g, 0.007 moles) in ethanol (40 ml) at reflux for 48 hours. The mixture was concentrated *in*  
*vacuo* and treated with saturated potassium carbonate. Extraction with chloroform (4x50

ml), drying and chromatography on silica gel eluting with 0-10% methanol/chloroform gave the title compound 1.0 g (69%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.25 (3H, t, J 8 Hz), 1.45 (6H, t, J 9 Hz), 2.08 (2H, s), 2.85 (6H, t, J 9 Hz), 4.05 (2H, q, J 8 Hz).

**Step 4. Quinuclidin-4-ylacetic acid hydrochloride**

- 5 Ethyl quinuclidin-4-ylacetate (1.0 g, 0.005 moles) was heated under reflux in 5M hydrochloric acid (60 ml) for 18 hours. Evaporation to dryness and trituration with acetone gave the title compound 0.93 g (89%) <sup>1</sup>H NMR (CD<sub>3</sub>SOCD<sub>3</sub>) 1.71 (6H, t, J 9 Hz), 2.15 (2H, s), 3.05 (6H, t, J 9 Hz), 10.35-10.55 (1H, br s), 12.19-12.29 (1H, br s).

**Step 5. Quinuclidin-4-ylacetyl chloride hydrochloride**

- 10 Quinuclidine-4-acetic acid hydrochloride (0.5 g, 0.0024 moles) was converted to the title compound using the method of Example 8, Step 4.

MS (+ve ion electrospray in methanol) m/z 183 (MH<sup>+</sup> for methyl ester, 100% showing complete conversion)

**Step 6. (3R)-3-Deoxy-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-(quinuclidin-4-yl)-acetate**

- 15 Quinuclidin-4-ylacetyl chloride hydrochloride (0.54 g, 0.0024 moles) and (3R)-3-deoxy-11-deoxy-3-methoxy-11-oxo-4-epimutilin (0.84 g, 0.0025 moles) were heated together in dry dimethylformamide (15 ml) at 100°C for 6 hours. The mixture was evaporated to dryness and the residue partitioned between saturated sodium hydrogen carbonate and  
20 chloroform. The organic layer was dried and chromatographed on silica gel eluting with 0-6% methanol/chloroform to give the title compound 0.4 g (39%) as a foam; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 0.87 (3H, d, J 7 Hz), 0.98 (3H, d, J 7 Hz), 1.05-1.70 (19 H, m), 1.95-2.03 (2H, m), 2.15 (2H, d, J 5 Hz), 2.17-2.21 (1H, m), 2.35-2.45 (1H, m), 2.85-2.97 (8H, m), 3.15 (3H, s), 3.35-3.45 (1H, m), 4.95 (1H, d, J 17 Hz), 5.30 (1H, d, J 12 Hz), 5.70 (1H, d, J 12  
25 Hz), 6.67 (1H, dd, J 17 Hz and J 10 Hz); MS (+ve ion electrospray) m/z 486 (MH<sup>+</sup>, 100%).

**Step 7. Mutilin 14-(quinuclidin-4-yl)-acetate**

- (3R)-3-Deoxy-11-deoxy-3-methoxy-11-oxo-4-epimutilin-14-(quinuclidin-4-yl)-acetate (0.37 g, 0.008 moles) in 1,4-dioxan (5 ml) was treated with concentrated hydrochloric acid  
30 (5 ml) and stirred at ambient temperature for 4 hours. Water (20 ml) was added and the mixture basified with sodium hydrogen carbonate. The product was extracted into chloroform (2 x 25 ml), dried (Na<sub>2</sub>SO<sub>4</sub>) filtered and evaporated to dryness to give the title compound as a white foam 0.33 g (92%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.7 (1H, d, J 7 Hz), 0.85 (1H, d, J 7 Hz), 1.1 (3H, s), 1.4 (3H, s), 2.85 (6H, t, J 9 Hz), 3.30-3.45 (1H, br s), 5.18 (1H, d, J 17 Hz), 5.31 (1H, d, J 10 Hz), 5.75 (1H, J, 10 Hz), 6.50 (1H, dd, J 17 and 10 Hz).  
35 MS (+ve ion electrospray) m/z 472 (MH<sup>+</sup>, 100%).

**Example 25 - Mutilin 14-(quinuclidin-4-ylmethyl)-aminoacetate**

**Step 1. 4-Cyanoquinuclidine**

Quinuclidin-4-ylcarbonylchloride hydrochloride (Example 8, Step 4) (3.4 g 0.016 moles) was dissolved in acetonitrile (150 ml) and treated with 35% ammonia solution (50 ml). The mixture was stirred for 18 hours at ambient temperature then concentrated to dryness *in vacuo*. 1 g of the residue was then treated with phosphorus oxychloride (8 ml) at reflux for 5 hours. The mixture was then concentrated *in vacuo* and the residue partitioned between saturated potassium carbonate and diethylether (4 x 50 ml). The combined organic extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. Column chromatography on silica gel eluting with 0-5% methanol/chloroform gave the title compound 0.34 g (75%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.85 (6H, t, J 10 Hz), 2.91 (6H, t, J 10 Hz)

10 **Step 2. 4- Aminomethylquinuclidine**

4- Cyanoquinuclidine (0.31 g, 0.0028 moles) was reduced with lithium aluminium hydride (0.45 g, 0.012 moles) in tetrahydrofuran (20 ml) at ambient temperature for 18 hours. Diethyl ether (20 mls) was added followed by water (1.8 ml) and 10%w/v aqueous sodium hydroxide (0.68 ml) and the mixture stirred for 30 minutes. The mixture was then filtered and the filtrate concentrated *in vacuo* to give the title compound 0.3 g (94%).

**Step 3. Mutilin 14-(quinuclidin-4-ylmethyl)-aminoacetate**

4-Aminomethylquinuclidine (0.2 g, 0.0014 moles) in chloroform (20 ml) was treated with diisopropylethylamine (0.54 g, 0.0042 moles) and mutilin 14-methane sulphonyloxyacetate (0.65 g, 0.0014 moles). The mixture was heated under reflux for 4 hours then allowed to cool. The solution was washed with saturated sodium hydrogen carbonate solution (2 x 20 ml). The organic phase was separated and dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. Chromatography on a Sep-Pak silica gel (10 g) cartridge eluting with 0-10% (9:1 methanol/35% ammonia solution) in chloroform gave the title compound 0.0065 g (1%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.71 (3H, d, J 7 Hz), 0.89 (3H, d, J 7 Hz), 1.1 (3H, s), 1.41 (3H, s), 2.80 (6H, t, J 10 Hz), 3.28 (2H, q, J 21 Hz), 5.20 (1H, d, J 17 Hz), 5.35 (1H, d, J 11 Hz), 5.75 (1H, d, J 8 Hz), 6.52 (1H, dd, J 17 and 11 Hz). MS (+ve ion electrospray) m/z 501 (MH<sup>+</sup>, 30%).

**Example 26 - Mutilin 14-[3-(quinuclidin-4-yl)-acrylate]**

30 **Step 1. N',O-Dimethylquinuclidin-4-yl amide**

Quinuclidin-4-ylcarbonylchloride hydrochloride (Example 8, Step 4) (16.5 g, 0.079 moles) in acetonitrile (600 ml) at 0°C was treated with N,O-Dimethylhydroxylamine hydrochloride (8.8g, 0.09 moles) and pyridine (20 ml, 0.24 moles) and stirred at ambient temperature for 18 hours. The mixture was concentrated *in vacuo* and the residue partitioned between saturated potassium carbonate and diethyl ether. The organics were dried (Na<sub>2</sub>SO<sub>4</sub>) filtered and evaporated to dryness to give the title compound 8.8 g (57%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.88 (6H, t, J 10 Hz), 2.91 (6H, t, J 10 Hz), 3.13 (3H, s), 3.65 (3H, s).

**Step 2. Quinuclidine-4-carboxaldehyde**

N,O-Dimethylquinuclidin-4-yl amide (8.77 g, 0.044 moles) in dry toluene at -70°C was treated with 1.5 molar diisobutylaluminiumhydride (45 ml, 0.067 moles) and allowed to warm to ambient temperature over 2 hours. The reaction was quenched with excess 5M hydrochloric acid, basified with potassium carbonate and extracted into diethyl ether. The organics were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. Chromatography on silica gel eluting with 0-10% (9:1 methanol/880 ammonia) in chloroform gave the title compound 1.3g (21%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.59 (6H, t, J 10 Hz), 2.90 (6H, t, J 10 Hz), 9.40 (1H, s).

**Step 3. Ethyl [3-(quinuclidin-4-yl)-acrylate]**

Triethylphosphonoacetate (1.6 ml, 0.0077 moles) in dimethoxyethane (50 ml) was treated with sodium hydride 60% dispersion in oil (0.35 g, 0.0088 moles) at ambient temperature for 1 hour. Quinuclidine-4-carboxaldehyde (1.0 g, 0.0072 moles) was then added and the mixture heated under reflux for 2 hours, allowed to cool and concentrated *in vacuo*. Chromatography of the residue on silica-gel, eluent as in Step 2, gave the title compound 0.71 g (47%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.29 (3H, t, J 10 Hz), 1.55 (6H, t, J 10 Hz), 2.99 (6H, t, J 10 Hz), 4.18 (2H, q, J 10 Hz), 5.65 (1H, d, J 19 Hz), 6.79 (1H, d, J 19 Hz).

**Step 4. 3-(Quinuclidin-4-yl)-acrylic acid hydrochloride**

Ethyl 3-(quinuclidin-4-yl)-acrylate (0.7 g, 0.0033 moles) was heated under reflux in 5 molar hydrochloric acid (30 ml) for 18 hours, cooled then concentrated *in vacuo* to an oil. Trituration with acetone gave the title compound as an off-white solid 0.43 g (60%). MS (+ve ion electrospray) m/z 182 (MH<sup>+</sup>, 100%).

**Step 5. 3-(Quinuclidin-4-yl)-acryloyl chloride hydrochloride**

The title compound was prepared from 3-(quinuclidin-4-yl)-acrylic acid as in the method of Example 8, Step 4 (0.24g, 100%). MS (+ve ion electrospray) m/z 196 (MH<sup>+</sup>, 100% - methyl ester from reaction with methanol).

**Step 6. (3R)-3-Deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin-14-[3'-(quinuclidin-4-yl)-acrylate]**

3-(Quinuclidin-4-yl)acryloyl chloride (0.24 g, 0.001 moles) and (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin (0.34 g, 0.001 moles) were heated together in dimethylformamide (15 ml) at 110°C for 18 hours. The mixture was allowed to cool and concentrated *in vacuo*. The residue was partitioned between chloroform and saturated sodium hydrogen carbonate solution. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to dryness. Chromatography on Sep-Pak silica gel 10 g cartridge eluting with 0-10% (9:1 methanol / 35% ammonia solution) in chloroform gave the title compound 0.035 g (6.5%); MS (+ve ion electrospray) m/z 498 (MH<sup>+</sup>, 100%).

**Step 7. Mutilin 14-[3-(quinuclidin-4-yl)-acrylate]**

The title compound was prepared from (3R)-3-Deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin-14-[3'-(quinuclidin-4-yl)-acrylate] (0.035 g, 0.00007 moles) as in the method of Example 24, Step 7 0.026 g, (76%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.61 (3H, d, J 7 Hz), 0.8 (3H, d, J 7 Hz), 1.1 (3H, s), 2.80 (6H, t, J 10 Hz), 5.12 (1H, d, J 17 Hz), 5.28 (1H, d, J 11



Hz), 5.49 (1H, d, J 15 Hz), 5.70 (1H, d, J 8 Hz), 6.49 (1H, dd, J 17 and 11 Hz), 6.64 (1H, d, J 15 Hz); MS (+ve ion electrospray) m/z 484 (MH<sup>+</sup>, 85%).

**Example 27 - Mutilin 14-[3-(Quinuclidin-4-yl)]-propionate**

5 **Step 1. 3-(Quinuclidin-4-yl)-propionic acid hydrochloride**

3-(Quinuclidin-4-yl)-acrylic acid (Example 26, Step 4) (0.2 g, 0.0009 moles) was hydrogenated at atmospheric pressure and ambient temperature over 10% palladium on charcoal (0.05 g) for 18 hours. The catalyst was filtered off and the filtrate evaporated to dryness to give the title compound 0.18 g (89%); MS (+ve ion electrospray) m/z 184  
10 (MH<sup>+</sup>, 100%).

**Step 2. 3-(Quinuclidin-4-yl)-propionylchloride hydrochloride**

The title compound was prepared from 3-(quinuclidin-4-yl)-propionic acid hydrochloride (0.18 g, 0.0008 moles) as in the method of Example 8, Step 4 0.19 g (100%). MS (+ve ion electrospray) m/z 198 (MH<sup>+</sup>, 100%)-methyl ester from reaction with methanol).

15 **Step 3. (3R)-3-Deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-[3'-(quinuclidin-4-yl)-propionate]**

The title compound was prepared from 3-(quinuclidin-4-yl)-propionyl chloride hydrochloride (0.19 g, 0.0008 moles) and (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin (0.27 g, 0.0008 moles) as in the method of Example 24, Step 6 0.19 g (48%).

20 MS (+ve ion electrospray) m/z 500 (MH<sup>+</sup>, 100%).

**Step 4. Mutilin 14-[3'-(quinuclidin-4-yl)-propionate]**

The title compound was prepared from (3R)-3-deoxo-11-deoxy-3-methoxy-11-oxo-4-epimutilin 14-[3'-(quinuclidin-4-yl)-propionate] (0.18 g, 0.0004 moles) as in the method of Example 24, Step 7 0.15 g (83%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.69 (3H, d, J 7 Hz), 0.87  
25 (3H, d, J 7 Hz), 1.15 (3H, s), 1.45 (3H, s), 2.85 (6H, t, J 10 Hz), 5.17 (1H, d, J 17 Hz), 5.33 (1H, d, J 11 Hz), 5.69 (1H, d, J 8 Hz), 6.51 (1H, dd, J 17 and 11 Hz). MS (+ve ion electrospray) m/z 486 (MH<sup>+</sup>, 100%).

**Example 28 - Mutilin 14-(quinuclidin-4-ylmethoxy)-acetate**

30 **Step 1. Quinuclidin-4-ylmethanol**

Quinuclidine-4-carboxylic acid hydrochloride (3.0 g, 0.016 moles) was treated with lithium aluminium hydride (2.5 g, 0.066 moles) in tetrahydrofuran (150 ml) at ambient temperature for 18 hours. The reaction was worked up as in the method of Example 25 Step 1 to give the title compound 2.24 g (100%). MS (+ve electrospray) m/z 142 (MH<sup>+</sup>, 100%).

35 **Step 2. Mutilin 14-(quinuclidin-4-ylmethoxy)-acetate**

Quinuclidin-4-ylmethanol (0.3 g, 0.002 moles) in dry dimethylformamide (5 ml) was treated with sodium hydride 60% dispersion in oil (0.095 g, 0.0022 moles) at ambient temperature for 1 hour. The mixture was then cooled to -10° C and mutilin 14-methanesulphonyloxyacetate (1.0 g, 0.002 moles) was added. The mixture was stirred for 4 hours

at ambient temperature then concentrated *in vacuo*. The residue was partitioned between saturated sodium hydrogen carbonate and chloroform. The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) filtered and evaporated to dryness. Chromatography on Sep-Pak silica gel (10 g) cartridge eluting with 0-10% (9:1 methanol/880 ammonia) in chloroform gave the title compound 0.12 g (12%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.71 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.15 (3H, s), 1.40 (3H, s), 2.85 (6H, t, J 10 Hz), 3.14 (2H, dd, J 10 and J 2.6 Hz), 3.93 (2H, q, J 17 Hz), 5.19 (1H, d, J 17 Hz), 5.35 (1H, d, J 11 Hz), 5.82 (1H, d, J 8 Hz), 6.52 (1H, dd, J 17 and J 11 Hz), 5.82 (1H, d, J 8 Hz), 6.52 (1H, dd, J 17 and J 11 Hz). MS (+ve ion electrospray)  $m/z$  502 ( $\text{MH}^+$ , 100%).

**Example 29 - Mutilin 14-[(3R)-quinuclidin-3-ylamino]-acetate**

The title compound was prepared from (R)-(+)-3-aminoquinuclidine dihydrochloride and mutilin 14-methanesulphonyloxyacetate as in the method of Example 28 Step 2 0.05 g (9%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.72 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.18 (3H, s), 1.45 (3H, s), 5.20 (1H, d, J 17 Hz), 5.35 (1H, d, J 11 Hz), 5.78 (1H, d, J 8 Hz), 6.52 (1H, dd, J 17 and J 11 Hz). MS (+ve ion electrospray)  $m/z$  487 ( $\text{MH}^+$ , 82%).

**Example 30 - Mutilin 14-(quinuclidin-4-yl-amino)-acetate**

**Step 1. 4-Aminoquinuclidine dihydrochloride**

Quinuclidin-4-ylcarbonylchloride (Example 8, Step 4) (1.0 g, 0.0048 moles) was treated with sodium azide (0.34 g, 0.005 moles) in dimethylformamide (10 ml) at 50°C for 18 hours. The mixture was concentrated *in vacuo* and the residue partitioned between saturated potassium carbonate and toluene. The toluene solution was separated, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and the filtrate was heated under reflux for 1 hour to give the isocyanate. The mixture was allowed to cool and then extracted with 5 M hydrochloric acid (3 x 20 ml). The combined acid extracts were then heated under reflux for 1 hour, cooled then evaporated to dryness. Trituration with acetone gave the title compound as a white solid 0.56 g (60%). M.S. (+ve ion electrospray)  $m/z$  127 ( $\text{MH}^+$ , 100%).

**Step 2. Mutilin 14-(quinuclidin-4-ylamino)-acetate**

The title compound was prepared from 4-aminoquinuclidine dihydrochloride and mutilin 14-methanesulphonyloxyacetate as in the method of Example 28 Step 2 0.023 g (3%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.7 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.17 (3H, s), 1.48 (3H, s), 2.95 (6H, t, J 10 Hz), 5.20 (1H, d, J 17 Hz), 5.35 (1H, d, J 11 Hz), 5.75 (1H, d, J 8 Hz), 6.49 (1H, dd, J 17 and J 11 Hz). M.S. (+ve ion electrsospray)  $m/z$  487 ( $\text{MH}^+$ , 100%).

**Example 31 - Mutilin 14-[4-(quinuclidin-4-yl)]-butyrate**

**Step 1. Quinuclidine-4-acetonitrile**

Quinuclidin-4-ylmethanol (1.94 g, 0.014 moles) was converted to the corresponding mesylate by treatment with methane sulphonyl chloride and triethylamine in chloroform.

The mesylate was dissolved in dimethylformamide (50 ml) and treated with sodium cyanide (1.4 g, 0.028 moles) at 120°C for 18 hours. The mixture was cooled and concentrated *in vacuo*. The residue was partitioned between saturated potassium carbonate and chloroform. The organic layer was separated and dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and

- 5 evaporated to dryness. Chromatography on silica gel eluting with 0-10% methanol/chloroform gave the title compound 1.5 g (72%). M.S. (+ve electrospray) m/z 151 (MH<sup>+</sup>, 100%).

#### Step 2. Quinuclidine-4-acetaldehyde

- 10 Quinuclidin-4-ylacetonitrile (3.0 g, 0.02 moles) in dry toluene (100 ml) was treated with 1.5 molar diisobutyl aluminium hydride (19.7 ml, 0.03 moles) at ambient temperature for 5 hours. The mixture was quenched by adding 2 M hydrochloric acid (50 ml) and stirring for 30 minutes. The mixture was then basified with potassium carbonate and extracted with chloroform. The organics were separated, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to dryness to give the title compound as an oil 2.2 g (72%). M.S. (+ve ion electrospray) m/z
- 15 154 (MH<sup>+</sup>, 100%).

#### Step 3. Mutilin 14-[4-(quinuclidin-4-yl)]-butyrate

- The title compound was prepared in 6 steps from quinuclidin-4-ylacetaldehyde analogously to Example 26 Steps 3-4 and Example 27 Steps 1-4 0.08 g (3% overall, 6 steps); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.65 (3H, d, J 7 Hz), 0.81 (3H, d, J 7 Hz), 1.10 (3H, s), 1.39 (3H, s),
- 20 2.95 (6H, t, J 10 Hz), 5.12 (1H, d, J 17 Hz), 5.27 (1H, d, J 11 Hz), 5.65 (1H, d, J 8 Hz), 6.43 (1H, dd, J 17 and J 11 Hz). M.S. (+ve electrospray) m/z 500 (MH<sup>+</sup>, 100%).

#### Example 32

##### (±) Mutilin 14-(1-azabicyclo[3,3,0]oct-4-ylmethylsulfanyl)-acetate

- 25 The title compound was prepared as in the method of Example 15 from (±)-1-azabicyclo[3,3,0]octan-4-ylmethanol (1.85 g, 0.007 moles) (Pizzorno, M.T., Albornico S.M., *J. Org. Chem.* (1974) **39**, 731). This gave 1.3 g (71%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.75 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.17 (3H, s), 1.45 (3H, s), 5.20 (1H, d, J 17 Hz), 5.35 (1H, d, J 11 Hz), 5.75 (1H, d, J 8 Hz), 6.50 (1H, dd, J 17 and 11 Hz). M.S. (+ve ion
- 30 electrospray) m/z 518 (MH<sup>+</sup>, 100%).

##### Example 33 - (±) Mutilin 14-(1-azabicyclo[3,3,0]oct-3-ylsulfanyl)-acetate

- The title compound was prepared as in the method of Example 15 from (±)-1-azabicyclo[3,3,0]octan-3-ol (0.6 g, 0.0047 moles) (Schnekenburger, *J. Pharm. Inst., Univ. Kiel*, Kiel, D-2300, Fed. Rep. Ger. Arch. Pharm. (1988), **321** (12), 925-9). This gave 0.21g
- 35 (9%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.72 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.18 (3H, s), 1.45 (3H, s), 5.20 (1H, d, J 17 Hz), 5.34 (1H, d, J 11 Hz), 5.74 (1H, d, J 8 Hz), 6.46 (1H, dd, J 17 and J 11 Hz). M.S. (+ve ion electrospray) m/z 504 (MH<sup>+</sup>, 35%).

**Example 34 - Mutilin 14-(endo 8-methyl-8-azabicyclo[3,2,1]oct-3-ylsulfanyl)-acetate**

The title compound was prepared as in the method of Example 15 from *exo* 8-methyl-8-azabicyclo[3,2,1]octan-3-ol (1.8 g, 0.0127 moles) (Nickon, A., Fieser, L.F., *J. American Chem. Soc.* (1952) 74, 5566). This gave 0.1 g (1.5%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.73 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.17 (3H, s), 1.47 (3H, s), 5.18 (1H, d, J 17 Hz), 5.32 (1H, d, J 11 Hz), 5.75 (1H, d, J 8 Hz), 6.47 (1H, dd, J 17 and 11 Hz). M.S. (+ve ion electrospray) m/z 518 (MH<sup>+</sup>, 100%).

**Example 35 - (±) Mutilin 14-(1-azabicyclo[4,3,0]non-4-ylsulfanyl)-acetate****Step 1. (±) 1-Azabicyclo[4,3,0]nonan-4-ol**

1-Azabicyclo[4,3,0]nonan-4-one (1.0 g, 0.0072 moles) (King, F.D., *J. Chem. Soc. Perkin. Trans. 1*, (1986) 447) in tetrahydrofuran (50 ml) was treated with lithium aluminium hydride (0.7 g, 0.0185 moles) at ambient temperature for 18 hours. Work up in the usual way gave the title compound 1.0 g (100%). M.S. (+ve ion electrospray) m/z 142 (MH<sup>+</sup>, 95%).

**Step 2. (±) Mutilin 14-(1-azabicyclo[4,3,0]non-4-ylsulfanyl)-acetate**

The title compound was prepared as in the method of Example 15 from (±) 1-Azabicyclo[4,3,0]nonan-4-ol (1.0 g, 0.0072 moles). This gave 1.12 g (28%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.72 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.20 (3H, s), 1.47 (3H, s), 5.21 (1H, d, J 17 Hz), 5.34 (1H, d, J 11 Hz), 5.77 (1H, d, J 8 Hz), 6.48 (1H, dd, J 17 and 11 Hz). M.S. (+ve ion electrospray) m/z 518 (MH<sup>+</sup>, 100%).

**Example 36 - (±) 19,20-Dihydromutilin 14-(1-azabicyclo[4,3,0]non-4-ylsulfanyl)-acetate**

The title compound was prepared as in the method of Example 15 from (±) 1-azabicyclo[4,3,0]nonan-4-ol (0.66 g, 0.0047 moles) and 19,20-dihydromutilin 14-methanesulphonyloxyacetate (2.43 g, 0.0047 mmoles) to give 0.44 g (18%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.71 (3H, d, J 7 Hz), 0.8 (3H, t, J 9 Hz), 1.45 (3H, s), 3.15 (2H, s), 5.65 (1H, d, J 8 Hz). M.S. (+ve ion electrospray) m/z 520 (MH<sup>+</sup>, 100%).

**Example 37 - Mutilin 14-(1-carboxymethylpiperidin-4-ylsulfanyl)-acetate****Step 1. tert-Butyl (piperidin-4-one-1-yl)-acetate**

4-Piperidone monohydrate hydrochloride (5 g, 0.033 moles) was treated with tert-butylbromoacetate (6.98 g, 0.037 moles) and potassium carbonate (13.65 g, 0.099 moles) in dimethylformamide (100 ml) at 100°C for 24 hours. The mixture was cooled and concentrated *in vacuo*. The residue was partitioned between saturated potassium carbonate solution and diethyl ether (2x50 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to dryness to give the title compound 7.36 g (94%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.45 (9H, s), 2.45 (4H, t, J 7 Hz), 2.3-2.4 (4H, m), 3.29 (2H, s).

**Step 2. tert-Butyl (piperidin-4-ol-1-yl)-acetate**

tert-Butyl (piperidin-4-one-1-yl)-acetate (3 g, 0.014 moles) was treated with sodium borohydride (1.13 g, 0.028 moles) in methanol (150 ml) at ambient temperature for 1 hour. Glacial acetic acid (1.68 g, 0.028 moles) was added and the mixture stirred for 15 minutes.

- 5 The mixture was concentrated *in vacuo* and the residue partitioned between saturated sodium carbonate and ethyl acetate. The organics were separated, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to dryness to give the title compound (2.9g, 96%). M.S. (+ve ion electrospray) m/z 216 (MH<sup>+</sup>, 100%).

**Step 3. Mutilin (1-carboxymethylpiperidin-4-ylsulfanyl)-acetate**

- 10 The title compound was prepared as in the method of Example 15 from tert-butyl(piperidin-4-ol-1-yl)-acetate (1.5 g, 0.007 moles). The tertbutyl ester group is hydrolysed in the workup. This gave 0.3 g (8%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.7 (3H, d, J 7 Hz), 0.88 (3H, d, J 7 Hz), 1.17 (3H, s), 1.47 (3H, s), 5.22 (1H, d, J 17 Hz), 5.35 (1H, d, J 11 Hz), 5.75 (1H, d, J 8 Hz), 6.45 (1H, dd, J 17 and 11 Hz). M.S. (+ve ion electrospray) m/z 536 (MH<sup>+</sup>, 100%).
- 15

**Example 38 - Mutilin 14-(piperidin-4-ylsulfanyl)-acetate****Step 1. 1(tert-Butoxycarbonyl)piperidin-4-ol**

- 1(tert-Butoxycarbonyl)-4-piperidone (5 g, 0.025 moles) was treated with sodium borohydride (1.89 g, 0.05 moles) as in the method of Example 37, Step 2 to give the title compound 5.07 g (100%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 1.45 (9 H, s), 1.29-1.41 (2 H, m), 2.42-3.05 (2H, m), 3.75-3.99 (3H, m).
- 20

**Step 2. Mutilin 14-(1-tert-butoxycarbonylpiperid-4-ylthio)-acetate**

- The title compound was prepared as in the method of Example 15 from 1-(tert-butoxycarbonyl)piperidin-4-ol (2.5 g, 0.012 moles). M.S. (-ve ion electrospray) m/z 576 (M-H, 100%).
- 25

**Step 3. Mutilin 14-(piperidin-4-ylsulfanyl)-acetate**

- The product from Step 2 was treated with trifluoroacetic acid (10 ml) in dichloromethane (100 ml) at 0°C for 2 hours. The mixture was concentrated *in vacuo* and the residue partitioned between saturated sodium hydrogen carbonate and chloroform. The organic layer was separated, dried (Na<sub>2</sub>SO<sub>4</sub>) filtered and evaporated to dryness. Chromatography on silica gel eluting with 0-10% (9:1 methanol/880 ammonia) in chloroform gave the title compound 1.01 g (26%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.75 (3H, d, J 7 Hz), 0.9 (3H, d, J 7 Hz), 1.18 (3H, s), 1.45 (3H, s), 5.20 (1H, d, J 17 Hz), 5.35 (1H, d, J 11 Hz), 5.80 (1H, d, J 8 Hz), 6.52 (1H, dd, J 17 and 11 Hz). M.S. (+ve ion electrospray) m/z 478 (MH<sup>+</sup>, 65%).
- 30
- 35

**Example 39 - Mutilin 14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate hydrochloride****Step 1. 1-Methyl-4-(hydroxymethyl)piperidine**

1-Methylpiperidine-4-carboxylic acid hydrochloride (*J. Med. Chem.* 1988, **31**, 812) (1 g, 0.007 mole) was added portionwise to a suspension of lithium aluminium hydride (1.3 g, 0.035 mole) in dry tetrahydrofuran (100 ml) under argon at 0°C. The mixture was heated under reflux overnight after which it was cooled to 0°C and treated dropwise with water (1.3 ml), 10% sodium hydroxide solution (1.95 ml) and water (3.25 ml) and stirred for 1 hour at room temperature. The resulting slurry was filtered through celite and the filtrate evaporated *in vacuo* to afford the title compound 0.90 g (99.7%) as a pale orange oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.18-1.53 (3H, m), 1.67-1.81 (2H, m), 1.83-2.12 (3H, m), 2.28 (3H, s), 2.79-2.94 (2H, m), 3.50 (2H, d, J 7 Hz); MS (+ve ion electrospray) m/z 130 (MH<sup>+</sup>).

**Step 2. (1-Methylpiperidin-4-ylmethylsulfanyl)-acetate**

Triphenylphosphine (3.67 g, 0.014 mole) was dissolved in dry tetrahydrofuran (25 ml) and cooled to 0°C under argon. Diisopropyl azodicarboxylate (2.75 ml, 0.014 mole) was added dropwise and the mixture was stirred at 0°C for 0.5 hour. The product of Step 1 (0.90 g, 0.007 mole) and thiolacetic acid (1.0 ml, 0.014 mole) in dry tetrahydrofuran (50 ml) were added dropwise and the mixture stirred at room temperature overnight. The solvent was removed *in vacuo* and the residue was partitioned between 1M hydrochloric acid and diethyl ether. The aqueous layer was washed with diethyl ether until all triphenylphosphine oxide had been removed, basified with solid potassium carbonate, extracted into dichloromethane, dried (magnesium sulfate) and evaporated *in vacuo* to afford the title compound 0.60 g (46%) as a pale yellow oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.22-1.59 (3H, m), 1.72-1.85 (2H, m), 1.95 (2H, dt, J 13 and 3 Hz), 2.28 (3H, s), 2.35 (3H, s), 2.80-2.98 (4H, m); MS (+ve ion electrospray) m/z 188 (MH<sup>+</sup>).

**Step 3. Mutilin 14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate hydrochloride**

The product of Step 2 (0.19 g, 0.001 mole) was dissolved in dry ethanol (10 ml) under argon and treated with sodium methoxide (0.054 g, 0.001 mole). The mixture was stirred for 1 hour and mutilin 14-methanesulfonyloxyacetate (0.456 g, 0.001 mole) was added. The mixture was stirred at room temperature overnight. The solvent was removed *in vacuo* and the residue partitioned between water and dichloromethane. The organic layer was dried (magnesium sulfate) and evaporated *in vacuo*. The residue was purified by column chromatography, eluting with dichloromethane to 15% methanol/ dichloromethane. The resulting gum was converted to the hydrochloride salt to afford the title compound 0.17 g (34%) as a white foam; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.73 (3H, d, J 7 Hz), 0.90 (3H, d, J 7 Hz), 5.23 (1H, dd, J 17 and 3 Hz), 5.35 (1H, dd, J 13 and 3 Hz), 5.73 (1H, d, J 7 Hz), 6.48 (1H, q, J 17 and 10 Hz), 12.26-12.69 (1H, br s); MS (+ve ion electrospray) m/z 506 (MH<sup>+</sup> free base).

**Example 40 - Mutilin 14-((3S,4R)-1-azabicyclo[2.2.1]hept-3-ylmethylsulfanyl)-acetate**  
**Step 1. (3S,4R)-1-Azabicyclo[2.2.1]hept-3-ylmethanol**

The title compound 0.60 g (84%) was prepared from (3S,4R)-1-azabicyclo[2.2.1]heptane-3-carboxylic acid hydrochloride (WO 98/05659, SmithKline Beecham) using the method of Example 1 Step 1; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.38-1.65 (1H, m), 1.83-2.00 (1H, m), 2.12-2.66 (7H, m), 2.78-3.05 (2H, m), 3.49-3.81 (2H, m); MS (+ve ion electrospray) m/z 128 (MH<sup>+</sup>).

**Step 2. [(3S, 4R)-1-Azabicyclo[2.2.1]hept-3-ylmethylsulfanyl]-acetate**

The title compound, 0.58g (66%) was prepared from the product of Step 1 using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.36-1.63 (2H, m), 1.90-2.01 (1H, m), 2.10-2.29 (1H, m), 2.34 (3H, s), 2.40-2.58 (4H, m), 2.78-2.96 (2H, m), 3.00-3.13 (2H, m); MS (+ve ion electrospray) m/z 186 (MH<sup>+</sup>).

**Step 3. Mutilin 14-[(3S,4R)-1-azabicyclo[2.2.1]hept-3-ylmethylsulfanyl]-acetate**

The title compound, 0.21 g (42%) was prepared from the product of Step 2 using the method of Example 1 Step 3. Purification of the compound was achieved by flash column chromatography on silica gel eluting with 10% methanol/dichloromethane; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.76 (3H, d, J 7 Hz), 0.90 (3H, d, J 7 Hz), 3.13 (2H, s), 5.20 (1H, dd, J 18 and 2 Hz), 5.34 (1H, dd, J 12 and 2 Hz), 5.78 (1H, d, J 7 Hz), 6.51 (1H, q, J 18 and 13 Hz); MS (+ve ion electrospray) m/z 504 (MH<sup>+</sup>).

**Example 41 - Mutilin 14-(quinuclidin-2-ylmethylsulfanyl)-acetate**

**Step 1. (Quinuclidin-2-ylmethylsulfanyl)-acetate**

The title compound, 0.78 g (55%) was prepared from quinuclidin-2-ylmethanol (*J. Am. Chem. Soc.*, 1988, **116**, 1278) using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.08-1.22 (1H, m), 1.40-1.58 (4H, m), 1.73-1.90 (2H, m), 2.35 (3H, s), 2.66-3.28 (7H, m); MS (+ve ion electrospray) m/z 158 (MH<sup>+</sup> thiol).

**Step 2. Mutilin 14-(quinuclidin-2-ylmethylsulfanyl)-acetate**

The title compound, 0.20 g (39%) was prepared from the product of Step 1 using the method of Example 1 Step 3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 1.75 (3H, d, J 7 Hz), 0.90 (3H, d, J 7 Hz), 3.18 (2H, d, J 7 Hz), 5.21 (1H, dd, J 18 and 2 Hz), 5.37 (1H, dd, J 12 and 2 Hz), 5.75 (1H, d, J 7 Hz), 6.50 (1H, q, J 18 and 12 Hz); MS (+ve ion electrospray) m/z 518 (MH<sup>+</sup>).

**Example 42 - Mutilin 14-(1-azabicyclo[2.2.1]hept-4-ylmethylsulfanyl)-acetate**

**Step 1. (1-Azabicyclo[2.2.1]hept-4-ylmethylsulfanyl)-acetate**

The title compound, 0.55 g (42%) was prepared from 1-azabicyclo[2.2.1]hept-4-yl methanol (WO 93/15080) using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.21-1.35 (2H, m), 1.50-1.68 (2H, m), 2.29 (2H, s), 2.38 (3H, s), 2.53-2.70 (2H, m), 2.99-3.05 (2H, m), 3.28 (2H, s); MS (+ve ion electrospray) m/z 186 (MH<sup>+</sup>).

**Step 2. Mutilin 14-(1-azabicyclo[2.2.1]hept-4-ylmethylsulfanyl)-acetate**

The title compound, 0.14 g (28%) was prepared from the product of Step 1 using the method of Example 1 Step 3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.78 (3H, d, J 7 Hz), 0.90 (3H, d, J 7 Hz), 3.16 (2H, s), 5.22 (1H, dd, J 18 and 2 Hz), 5.37 (1H, dd, J 12 and 2 Hz), 5.78 (1H, d, J 8 Hz), 6.50 (1H, q, J 18 and 12 Hz); MS (+ve ion electrospray) m/z 504 (MH<sup>+</sup>).

5

**Example 43 - Mutilin 14-((3R,4S)-1-azabicyclo[2.2.1]hept-3-ylmethylsulfanyl)-acetate**  
**Step 1. (3R, 4S)-1-Azabicyclo[2.2.1]hept-3-yl methanol**

The title compound, 0.68 g (95%) was prepared from (3R,4S)-1-azabicyclo[2.2.1] heptane-carboxylic acid using the method of Example 1 Step 1; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.37-1.71 (2H, m), 1.82-2.00 (1H, m), 2.10-2.72 (6H, m), 2.77-3.05 (2H, m), 3.47-3.76 (2H, m); MS (+ve ion electrospray) m/z 128 (MH<sup>+</sup>).

10

**Step 2. [(3R,4S)-1-Azabicyclo[2.2.1]hept-3-ylmethylsulfanyl]-acetate**

The title compound, 0.22 g (25%) was prepared from the product of Step 1 using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.40-1.70 (2H, m), 1.93-2.09 (1H, m), 2.12-2.31 (1H, m), 2.35 (3H, s), 2.51-2.70 (4H, m), 2.78-2.98 (2H, m), 3.0-3.15 (2H, m); MS (+ve ion electrospray) m/z 186 (MH<sup>+</sup>).

15

**Step 3. Mutilin 14-((3R,4S)-1-azabicyclo[2.2.1]hept-3-ylmethylsulfanyl)-acetate**

The title compound 0.12 g (20%) was prepared from the product of Step 2 using the method of Example 1 Step 3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.72 (3H, d, J 7 Hz), 0.89 (3H, d, J 7 Hz), 3.13 (2H, s), 5.21 (1H, dd, J 18 and 2 Hz), 5.35 (1H, dd, J 12 and 2 Hz), 5.76 (1H, d, J 7 Hz), 6.50 (1H, q, J 18 and 12 Hz); MS (+ve ion electrospray) m/z 504 (MH<sup>+</sup>).

20

**Example 44 - Mutilin 14-(1-azabicyclo[3.2.1]oct-5-ylmethylsulfanyl)-acetate**

**Step 1. 1-Azabicyclo[3.2.1]oct-5-ylmethanol**

The title compound, 2.05 g (93%) was prepared from 1-azabicyclo[3.2.1]octane-5-carboxylic acid hydrochloride (*J. Med. Chem.*, 1991, 34, 2726-2735) using the method of Example 1 Step 1; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.39-1.90 (5H, m), 2.61 (2H, s), 2.70 (4H, m), 3.35 - 3.75 (4H, m); MS (+ve ion electrospray) m/z 142 (MH<sup>+</sup>).

25

**Step 2. [1-Azabicyclo[3.2.1]oct-5-ylmethylsulfanyl]-acetate**

The title compound, 1.0 g (35%) was prepared from the product of Step 1 using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.45-1.89 (6H, m), 2.47 (3H, s), 2.60 (2H, s), 2.70-2.94 (3H, m), 3.00-3.17 (3H, m); MS (+ve ion electrospray) m/z 200 (MH<sup>+</sup>).

30

**Step 3. Mutilin 14-(1-azabicyclo[3.2.1]oct-5-ylmethylsulfanyl)-acetate**

The title compound, 0.19 g (7%) was prepared from the product of Step 2 using the method of Example 1 Step 3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.73 (3H, d, J 7 Hz), 0.90 (3H, d, J 7 Hz), 5.20 (1H, dd, J 18 and 2 Hz), 5.37 (1H, dd, J 12 and 2 Hz), 5.76 (1H, d, J 7 Hz), 6.48 (1H, q, J 18 and 12 Hz); MS (+ve ion electrospray) m/z 518 (MH<sup>+</sup>).

35



**Example 45 - Mutilin 14-[(R)-1-Methylpiperid-2-ylmethylsulfanyl]-acetate****Step 1. (R)-1-Ethylcarbamoylpiperidine-2-carboxylic acid**

L-Pipecolinic acid (0.50 g, 0.004 mole) in dry dichloromethane (10 ml) was cooled to 0°C under argon and treated with triethylamine (0.65 ml, 0.0046 mole) followed dropwise by ethyl chloroformate (0.37 ml, 0.004 mole) in dry dichloromethane (2 ml). The mixture was stirred overnight at room temperature after which it was diluted with dichloromethane, washed with 5M hydrochloric acid and the organic layer dried (magnesium sulfate) and evaporated *in vacuo* to afford the title compound 0.60 g (77%) as an orange oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.12-1.84 (8H, m), 2.15-2.40 (1H, m), 2.88-3.20 (1H, m), 3.90-4.28 (3H, m), 4.77-5.07 (1H, m), 5.68-6.82 (1H, br s).

**Step 2. (R)-1-Methylpiperidin-2-ylmethanol**

The product of Step 1 (0.60 g, 0.003 mole) in dry tetrahydrofuran (10 ml) was added dropwise to a suspension of lithium aluminium hydride (0.57 g, 0.015 mole) in dry tetrahydrofuran (20 ml). The mixture was heated under reflux for 2 hours and stirred at room temperature overnight. The reaction mixture was cooled to 0°C and water (0.5 ml) was added dropwise, followed by 10% sodium hydroxide solution (0.9 ml) and water (1.4 ml). The mixture was stirred for 1 hour, filtered through Celite and the filtrate evaporated *in vacuo* to afford the title compound 0.31 g (80%) as a pale yellow oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.17-2.00 (8H, m), 2.14 (1H, dt, J 13 and 2 Hz), 2.30 (3H, s), 2.76-2.96 (1H, m), 3.40 (1H, dd, J 13 and 1 Hz), 3.88 (1H, dd, J 12 and 5 Hz); MS (+ve ion electrospray) m/z 130 (MH<sup>+</sup>).

**Step 3. [(R)-1-Methylpiperid-2-ylmethylsulfanyl]-acetate**

The title compound, 0.30 g (71%) was prepared from the product of Step 2 using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.16-1.76 (6H, m), 2.00-2.18 (2H, m), 2.29 (3H, s), 2.35 (3H, s), 2.80-2.92 (1H, m), 3.00-3.23 (2H, m).

**Step 4. Mutilin 14-[(R)-1-Methylpiperid-2-ylmethylsulfanyl]-acetate**

The title compound, 0.19 g (22%) was prepared from the product of Step 3 using the method of Example 1 Step 3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.75 (3H, d, J 7 Hz), 0.89 (3H, d, J 7 Hz), 3.11 (2H, s), 3.36 (1H, q, J 12 and 7 Hz), 5.19 (1H, dd, J 18 and 2 Hz), 5.35 (1H, dd, J 12 and 2 Hz), 5.75 (1H, d, J 7 Hz), 6.50 (1H, q, J 18 and 12 Hz); MS (+ve ion electrospray) m/z 506 (MH<sup>+</sup>).

**Example 46 - Mutilin 14-[(S)-1-Methylpyrrolid-2-ylmethylsulfanyl]-acetate****Step 1. [(S)-1-Methylpyrrolid-2-ylmethylsulfanyl]-acetate**

The title compound, 0.64 g (85%) was prepared from (S)(-)-1-methyl-2-pyrrolid-2-ylmethanol using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.44-1.61 (1H, m), 1.65-1.85 (2H, m), 1.87-2.04 (1H, m), 2.15-2.42 (2H, m), 2.35 (3H, s), 2.38 (3H, s), 2.82-

2.94 (1H, m), 3.05-3.14 (1H, m), 3.28 (1H, dd, J 13 and 3 Hz); MS (-ve ion electrospray) m/z 130 (M-H for thiol).

**Step 2. Mutilin 14-[(S)-1-Methylpyrrolid-2-ylmethylsulfanyl]-acetate**

The title compound, 0.17 g (23%) was prepared from the product of Step 1 using the method of Example 1 Step 3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.76 (3H, d, J 7 Hz), 0.90 (3H, d, J 7 Hz), 3.18 (2H, s), 3.35 (1H, q, J 10 and 7 Hz), 5.20 (1H, dd, J 18 and 2 Hz), 5.35 (1H, dd, J 12 and 2 Hz), 5.75 (1H, d, J 7 Hz), 6.50 (1H, q, J 18 and 12 Hz); MS (+ve ion electrospray) m/z 492 (MH<sup>+</sup>).

**Example 47 - Mutilin 14-[(R)-1-methylpiperid-3-ylmethylsulfanyl]-acetate**

**Step 1. (R)-Ethyl 1-ethylcarbamoylpiperidine-3-carboxylate**

(R)-Ethyl nipecotate (*J. Org. Chem.*, **56**, 1991, 1166-1170) (3.0 g, 0.019 mole) in dry dichloromethane (50 ml) was cooled to 0°C under argon. Triethylamine (3.19 ml, 0.023 mole) was added, followed dropwise by ethyl chloroformate (1.83 ml, 0.019 mole) in dry dichloromethane (6 ml) and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with dichloromethane, washed with water, dried (magnesium sulfate) and evaporated *in vacuo* to afford the title compound 3.45 g (79%) as a colourless oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 1.28 (6H, t, J 7 Hz), 1.38-1.82 (3H, m), 2.00-2.15 (1H, m), 2.38-2.55 (1H, m), 2.77-3.13 (2H, m), 3.91-4.04 (1H, m), 4.07-4.35 (5H, m)

**Step 2. (R)-1-Methylpiperid-3-ylmethanol**

The title compound, 1.8 g (92%) was prepared from the product of Step 1 using the method of Example 7 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 0.90-1.12 (1H, m), 1.50-1.90 (5H, m), 1.94-1.99 (1H, m), 2.25 (3H, s), 2.57-2.74 (1H, m), 2.79-2.92 (1H, m), 3.14-3.71 (3H, m); MS (+ve ion electrospray) m/z 130 (MH<sup>+</sup>).

**Step 3. [(R)-1-Methylpiperid-3-ylmethylsulfanyl]-acetate**

The title compound, 0.59 g (81%) was prepared from the product of Step 2 using the method of Example 1 Step 2; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 0.87-1.06 (1H, m), 1.44-1.94 (6H, m), 2.27 (3H, s), 2.34 (3H, s), 2.62-2.92 (4H, m); MS (+ve ion electrospray) m/z 188 (MH<sup>+</sup>)

**Step 4. Mutilin 14-[(R)-1-methylpiperid-3-ylmethylsulfanyl]-acetate**

The title compound, 0.26 g (34%) was prepared from the product of Step 3 using the method of Example 1 Step 3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.74 (3H, d, J 7 Hz), 0.90 (3H, d, J 7 Hz), 3.12 (2H, d, J 2 Hz), 3.30-3.44 (1H, m), 5.22 (1H, dd, J 18 and 2 Hz), 5.38 (1H, dd, J 12 and 2 Hz), 5.76 (1H, d, J 7 Hz), 6.50 (1H, q, J 18 and 12 Hz); MS (+ve ion electrospray) m/z 506 (MH<sup>+</sup>).

**Example 48 - Mutilin 14-(quinuclidin-4-ylmethylsulfanyl)-acetate**

A solution of triphenylphosphine (1.1 g, 0.0042 mole) in dry tetrahydrofuran (50 ml) was ice-cooled under argon and treated with diisopropyl azodicarboxylate (0.85 g, 0.0042

mole). After 30 minutes a solution of thiolacetic acid (0.315 ml, 0.0042 mole) and quinuclidin-4-ylmethanol (0.565 g, 0.0042 mole) in dry tetrahydrofuran was added dropwise. The mixture was then allowed to stand at 5°C for 72 hours. Following concentration *in vacuo* the residue was partitioned between diethyl ether and 1M  
 5 hydrochloric acid. The aqueous phase was washed with diethyl ether then concentrated *in vacuo* to give a solid (0.65 g). The solid was dissolved in ethanol and treated with potassium tert-butoxide (0.785 g, 0.007 mole). After stirring for 30 minutes mutilin 14-methanesulfonyloxyacetate (1.38 g, 0.003 mole) was added. The mixture was stirred under argon for 18 hours then concentrated *in vacuo*. The residue was partitioned between  
 10 chloroform and water. The organic phase was washed with brine, dried over magnesium sulfate and concentrated *in vacuo*. Chromatography on silica gel eluting with chloroform / methanol / 35% ammonia solution (10/1/0.1) provided the title compound 0.478 g (31%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.74 (3H, d, J 6.5 Hz), 0.88 (3H, d, J 6.7 Hz), 1.17 (3H, s), 1.40 (6H, t, J 8 Hz), 1.49 (3H, s), 2.47 (2H, s), 2.87 (6H, t, J 8 Hz), 3.0 (2H, s), 3.36 (1H, m), 5.1 to 5.4 (2H, m), 5.75 (1H, d, J 8.3 Hz), 6.48 (1H, m); MS (+ve ion electrospray) m/z  
 15 518 (MH<sup>+</sup>, 100%).

**Example 49 - Mutilin 14-(8-methyl-8-azabicyclo [3.2.1]oct-3-ylmethylsulfanyl)-acetate**  
**Step 1. (8-methyl-8-azabicyclo[3.2.1]oct-3-yl)methanol**

20 The title compound was prepared from 8-methyl-8-azabicyclo[3.2.1]octane-3-carboxylic acid hydrochloride salt (WO 98/05659 Example 25, Step 3) using the process described in Example 24 Step 1 0.78 g (100%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 1.3-2.0 (9H, m), 2.25 (3H, s), 3.16 (2H, m), 3.44 (2H, d, J 6.3 Hz); MS (+ve ion electrospray) m/z 156 (MH<sup>+</sup>, 100%).

**Step 2. Mutilin 14-(8-methyl-8-azabicyclo[3.2.1]oct-3-ylmethylsulfanyl)-acetate**

25 The title compound was prepared from (8-methyl-8-azabicyclo[3.2.1]oct-3-yl)methanol and mutilin 14-methanesulfonyloxyacetate using the process described in Example 15 0.101 g (19%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.74 (3H, d, J 6.5 Hz), 0.88 (3H, d, J 7.0 Hz), 1.25 (3H, s), 1.49 (3H, s), 2.34 (3H, s), 2.48 (2H, d), 3.1 (2H, s), 3.15 (2H, m), 3.36 (1H, m), 5.1-5.4 (2H, m), 5.74 (1H, d, J 8.5 Hz), 6.48 (1H, m); MS (+ve ion electrospray) m/z  
 30 533 (MH<sup>+</sup>, 85%).

**Example 50 - Mutilin 14-(*exo*-8-methyl-8-azabicyclo[3.2.1]oct-3-ylsulfanyl)-acetate**

The title compound was prepared from *endo*-8-methyl-8-azabicyclo[3.2.1]octan-3-ol and mutilin 14-methanesulfonyloxyacetate using the process described in Example 15 0.09 g  
 35 (17%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.74 (3H, d, J 6.7 Hz), 0.99 (3H, d, J 7.5 Hz), 1.18 (3H, s), 1.63 (3H, s), 2.28 (3H, s), 3.0 (1H, m), 3.13 (2H, s), 3.16 (2H, m), 3.36 (1H, m), 5.15 to 5.37 (2H, m), 5.77 (1H, d, J 8.3 Hz), 6.49 (1H, m); MS (+ve ion electrospray) m/z 518 (MH<sup>+</sup>, 100%).

**Example 51 - Mutilin 14-[(3-(quinuclidin-4-ylsulfanyl))-propionate****Step 1. Mutilin 14-acrylate-11-trifluoroacetate**

Mutilin 11-trifluoroacetate (WO 97/25309 Example 85 step 2) (3.0 g, 0.0072 mole),  
 5 triethylamine (3.74 g, 0.037 mole) and a catalytic amount of 4-dimethylaminopyridine in dichloromethane (100 mls) was treated with acryloyl chloride (3.33 g, 0.037 mole) overnight at room temperature under argon. The reaction mixture was partitioned between water and dichloromethane. The organic layer was dried over magnesium sulfate and the solvents removed *in vacuo*. Chromatography of the residue on silica gel eluting with  
 10 ethylacetate / petroleum ether 40-60° (1:10) provided the title compound 1.25 g (37%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.69 (3H, d, J 6.6 Hz), 0.84 (3H, d, J 7 Hz), 1.06 (3H, s), 1.52 (3H, s), 2.1 to 2.4 (4H, m), 2.65 (1H, m), 5.0 (1H, d, 6.9 Hz), 5.20-5.37 (2H, m), 5.72-5.86 (2H, m), 6.0-6.1 (1H, m), 6.3-6.5 (2H, m).

**Step 2. Mutilin 14-[(3-(quinuclidin-4-ylsulfanyl))-propionate**

15 Mutilin 14-acrylate-11-trifluoroacetate (0.376 g, 0.008 mole) was treated with prepared potassium quinuclidin-4-sulfanate from quinuclidin-4-thiol hydrochloride (0.145 g, 0.0008 mole) and potassium tert-butoxide (0.094 g, 0.000838 mole) in ethanol (15 ml) under argon at room temperature overnight. Solvents were removed *in vacuo* and the residue chromatographed on silica gel using chloroform/methanol/35% ammonia solution  
 20 (10:1:0.1) mixture. This chromatographed product (0.262 g) was dissolved in tetrahydrofuran/water (5:1) (6 ml) and treated with 0.5 M sodium hydroxide solution (1 ml) for 3 hours at room temperature. The reaction mixture was concentrated *in vacuo*. The residue was chromatographed on silica gel using chloroform/methanol/35% ammonia solution (9:1:0.1) mixture to provide the title compound 0.152 g (36% overall);  
 25 <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.72 (3H, d, J 6.5 Hz), 0.87 (3H, d, J 7.0 Hz), 1.08 (3H, s), 1.63 (3H, s), 1.69 (6H, t, J 8 Hz), 2.00-2.47 (7H, m), 2.74 (2H, t, J 7.8 Hz), 3.36 (1H, m), 5.17-5.38 (2H, m), 5.74 (1H, d, J 8.5 Hz), 6.52 (1H, m); MS (+ve ion electrospray) m/z 518 (MH<sup>+</sup>, 100%).

**Example 52 - Mutilin 14-[3-(quinuclidin-4-ylmethylsulfanyl))-propionate**

30 To an ice-cooled solution of triphenylphosphine (1.1 g, 0.0042 mole) in dry tetrahydrofuran (50 ml) was added dropwise diisopropyl azodicarboxylate (0.85 g, 0.0042 mole). After 30 mins a solution of thiolacetic acid (0.335 g, 0.0042 mole) and quinuclidin-4-ylmethanol (Example 28, Step 1) (0.565 g, 0.004 mole) in dry tetrahydrofuran (20 ml) was added  
 35 dropwise. The mixture was stirred under argon for 72 hours, evaporated *in vacuo* and taken up in ether. The ethereal solution was extracted with 1M hydrochloric acid. The aqueous extract was washed with ether and evaporated to dryness to give a solid (0.65 g). The title compound was prepared from the latter solid and mutilin 14-acrylate-11-trifluoroacetate (Example 51, Step 1) according to the procedure of Example 51, Step 2

0.41 g (80%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.72 (3H, d, J 6.8 Hz), 0.87 (3H, d, J 7 Hz), 1.09 (3H, s), 1.45 (3H, s), 1.48 (6H, t, 8 Hz), 2.46 (2H, s), 2.52 (2H, m), 2.75 (2H, m), 2.95 (6H, t, J 7.8 Hz), 3.44 (1H, m), 5.28 (2H, m), 5.75 (1H, d, J 8.5 Hz), 6.52 (1H, m); MS (+ve ion electrospray)  $m/z$  532 ( $\text{MH}^+$ , 100%).

5

**Example 53 - Mutilin 14-[3-(1-methylpiperid-4-ylsulfanyl)]-propionate**

A solution of triphenylphosphine (5.51 g, 0.021 mole) in dry tetrahydrofuran (100 ml) treated with diisopropyl azodicarboxylate (4.25 g, 0.021 mole). After 30 minutes a solution of 1-methylpiperidin-4-ol (2.3 g, 0.02 mole) and thiolacetic acid (1.54 g, 0.02 mole) in dry tetrahydrofuran (50 ml) was added over a period of 30 minutes. The mixture was stirred overnight, concentrated in vacuo and the residue taken up in ether. The ethereal solution was extracted with 1M hydrochloric acid. The aqueous extract was washed with ether, evaporated to dryness and dried *in vacuo* to give a yellow gum (2.4 g). A portion of the gum (0.252 g) was treated with sodium methoxide (0.120 g) in ethanol and subsequently with mutilin 14-acrylate-11-trifluoroacetate (Example 51, Step 1) (0.376 g) according to the procedure of Example 51, Step 2 to give the title compound 0.3 g (74%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.73 (3H, d, J 6.8 Hz), 0.87 (3H, d, J 7.0 Hz), 1.17 (3H, s), 1.46 (3H, s), 2.18 (2H, m), 2.25 (3H, s), 2.40 (2H, m), 2.51 (1H, m), 2.80 (4H, m), 3.35 (1H, m), 5.27 (2H, m), 5.74 (1H, d, 8.3 Hz), 6.52 (1H, m); MS (+ve ion electrospray)  $m/z$  506 ( $\text{MH}^+$ , 100%).

20

**Example 54 - 19,20-Dihydromutilin 14-(1-methylpiperid-4-ylsulfanyl)-acetate**

**Step 1. 19,20-Dihydromutilin 14-methanesulfanyloxyacetate**

The title compound was prepared from 19,20-dihydropleuromutilin (A. Birch *et al*, *Tetrahedron* (1966) Suppl. 8 part II, 359-387) using the literature process for pleuromutilin (H. Egger and H. Reinshagen, *J. Antibiotics* 29 (9), 915);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.71 (3H, d, J 7 Hz), 0.77 (3H, t, 7.5 Hz), 0.95 (3H, d, J 8.5 Hz), 0.97 (3H, s), 1.42 (3H, s), 3.21 (3H, s), 3.42 (1H, m), 4.66 (2H, m), 5.72 (1H, d, 8.2 Hz).

**Step 2. 19,20-Dihydromutilin 14-(1-methylpiperidin-4-ylsulfanyl)-acetate**

The title compound was prepared from 4-hydroxy-1-methylpiperidine and 19,20-dihydromutilin 14-methanesulfonyloxyacetate using the process described in Example 15 0.42 g (83%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) *inter alia* 0.71 (3H, d, J 6.8 Hz), 0.78 (3H, t, J 7.6 Hz), 0.94 (3H, d, J 7.6 Hz), 0.97 (3H, s), 1.43 (3H, s), 2.25 (3H, s), 2.42 (1H, m), 2.81 (2H, m), 3.42 (1H, t, J 6 Hz), 5.63 (1H, d, J 8 Hz); MS (+ve ion electrospray)  $m/z$  494 ( $\text{MH}^+$ , 75%).

35

**Example 55 - 19,20-Dihydromutilin 14-(8-methyl-8-azabicyclo[3.2.1]oct-3-ylmethylsulfanyl)-acetate**

The title compound was prepared from 19,20-dihydromutilin 14-methanesulfonyloxyacetate (Example 54, step 1) and 8-methyl-8-azabicyclo[3.2.1]oct-3-ylmethanol using the process described in Example 15, 0.335 g (45%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.71 (3H, d, J 6.5 Hz), 0.79 (3H, t, J 7.3 Hz), 0.93 (3H, d, J 7.0 Hz), 0.97 (3H, s), 1.0 to 2.2 (27H, m), 2.28 (3H, s), 2.41 (1H, m), 3.11 (2H, s), 3.17 (2H, m), 3.42 (1H, m), 5.62 (1H, d, J 8.3 Hz); MS (+ve ion electrospray) m/z 520 (MH<sup>+</sup>, 60%).

#### Example 56 - Mutilin 14-[4-(quinuclidin-4-ylsulfanyl)]-butyrate

##### Step1. Mutilin 14-(4-bromobutyrate)-11-trifluoroacetate

Mutilin 11-trifluoroacetate (WO 97/25309, Example 85, Step 2) (1.25 g, 0.003 mole) and pyridine (0.237 g, 0.003 mole) in dry dichloromethane (20 ml) were treated with 4-bromobutyroyl chloride (0.56 g, 0.003 mole) for 72 hours. The mixture was concentrated *in vacuo* and the resulting residue chromatographed on silica gel using dichloromethane, providing the title compound, 1.5 g (93%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.72 (3H, d, J 6.7 Hz), 0.83 (3H, d, J 7 Hz), 1.05 (3H, s), 1.43 (3H, s), 2.62 (1H, t, J 7 Hz), 3.46 (2H, t, J 6 Hz), 5.0 (1H, d, J 6.7 Hz), 5.3 (2H, m), 5.69 (1H, d, J 8 Hz), 6.37 (1H, m); MS (+ve ion electrospray) m/z 532 (MH<sup>+</sup>, 40%).

##### Step2. Mutilin 14-[4-(quinuclidin-4-ylsulfanyl)]-butyrate

Quinuclidin-4-thiol hydrochloride (0.359 g, 0.002 mole) in ethanol (10 ml) was treated with sodium methoxide (0.216 g, 0.004 mole). After 30 minutes mutilin 14-(4-bromobutyrate) (0.565 g, 0.001 mole) was added and the mixture allowed to stand overnight under argon. The reaction mixture was concentrated *in vacuo* and the residue partitioned between water and chloroform. The organic layer was dried over magnesium sulfate and concentrated *in vacuo*. The residue was chromatographed on silica gel using chloroform/methanol/35% ammonia solution (10:1:0.1) to give title compound, 0.190 g (35%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.71 (3H, d, J 6.5 Hz), 0.87 (3H, d, J 6.8 Hz), 1.16 (3H, s), 1.59 (3H, s), 1.81 (14H, m), 2.06 (2H, t, J 8.5 Hz), 2.49 (2H, t, J 7.3 Hz), 2.94 (6H, t, J 7.3 Hz), 3.35 (1H, m), 5.29 (2H, m), 5.75 (1H, d, J 8.5 Hz), 6.52 (1H, m); MS (+ve ion electrospray) m/z 532 (MH<sup>+</sup>, 100%).

#### Example 57

##### 1,2-Didehydromutilin 14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate

##### Step 1. 1,2-Didehydromutilin 11-dichloroacetate

A solution of 1,2-didehydromutilin (1.41 g, 0.0044 mole) (prepared by analogy to the procedure described for 1,2-didehydropleuromutilin, G. Schulz and H. Berner, *Tetrahedron*, 1984, 40, 905-17), pyridine (0.56 ml, 0.0066 mole) and N,N-dimethylaminopyridine (0.02 g) in tetrahydrofuran (30 ml) was treated with dichloroacetic anhydride (1.16 g, 0.0048 mole) in tetrahydrofuran (5ml). After 18 hours the mixture was concentrated *in vacuo* and the residue partitioned between ethyl acetate and dilute hydrochloric acid. The organic

phase was separated, washed with water and brine, dried over magnesium sulphate and the solvent removed in vacuo. Chromatography of the residue on silica gel eluting with 20% ethyl acetate in hexanes gave the title compound (1.3 g, 69%) as a colourless solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 4.33 (1H, d, J 7.7Hz), 4.57 (1H, d, J 7.0Hz), 5.34 (1H, d, J 11.2Hz), 5.48 (1H, d, J 17.8Hz), 5.99 (1H, s), 6.10 (1H, d, J 6.1Hz), 6.11 (1H, dd, J 17.8 and 11.2 Hz), 7.67 (1H, d, J 6.1 Hz).

**Step 2. 1,2-Didehydromutilin 11 dichloroacetate-14-chloroacetate**

A solution of 1,2-didehydromutilin 11 dichloroacetate (1.2 g, 0.0028 mole), pyridine (0.7 ml) and N,N-dimethylaminopyridine (0.01 g) in dichloromethane (10 ml) at 0°C was treated with chloroacetyl chloride (0.33 ml, 0.0042 mole). After stirring at room temperature for 18 hours the mixture was concentrated in vacuo and the residue partitioned between ethyl acetate and dilute hydrochloric acid. The organic phase was separated, washed with water and brine, dried over magnesium sulphate and the solvent removed in vacuo. Chromatography of the residue on silica gel eluting with 20% ethyl acetate in hexanes gave the title compound (0.7 g, 50%) as a colourless solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.79 (3H, d, J 6.8Hz), 1.04 (3H, d, J 7.1Hz), 1.10 (3H, s), 1.58 (3H, s), 4.00 (2H, s), 4.60 (1H, d, J 7.0Hz), 5.30 (1H, d, J 17.7Hz) 5.36 (1H, d, J 11.7Hz), 5.70 (1H, d, J 8.6Hz), 5.97 (1H, s), 6.10 (1H, d, J 6.2Hz), 6.34 (1H, dd, J 17.7 and 11.7Hz), 7.66 (1H, d, J 6.2Hz).

**Step 3. 1,2-Didehydromutilin 11-dichloroacetate-14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate**

The title compound (0.36 g, 49%) was prepared from 1,2-didehydromutilin 11-dichloroacetate-14-chloroacetate (0.7 g, 0.0012 mole) and (1-methylpiperidin-4-ylmethylsulfanyl)-acetate (0.224 g, 0.0012 mole) using the process described in Example 39 Step 3. <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.80 (3H, d, J 6.3Hz), 1.03 (3H, d, J 7.0Hz), 1.09 (3H, s), 1.56 (3H, s), 2.26 (3H, s), 3.13 (2H, s), 4.60 (1H, d, J 6.8Hz), 5.30 (1H, d, J 17.5Hz), 5.34 (1H, d, J 10.7Hz), 5.66 (1H, d, J 8.4Hz), 5.97 (1H, s), 6.09 (1H, d, J 6.1Hz), 6.34 (1H, dd, J 17.5 and 10.7Hz), 7.65 (1H, d, J 6.1Hz); MS (+ve ion electrospray) 616 and 614 (MH<sup>+</sup>).

**Step 4. 1,2-Didehydromutilin 14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate**

A solution of 1,2-didehydromutilin 11-dichloroacetate-14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate (0.18 g, 0.0003 mole) in dioxane (3 ml) was treated with an aqueous potassium hydroxide (1M, 0.36ml). After stirring at room temperature for 1 hour the mixture was neutralised with dilute hydrochloric acid and the solvent evaporated in vacuo. The residue was partitioned between ethyl acetate and sodium hydrogen carbonate solution. The organic phase was separated, washed with water and brine, dried over magnesium sulphate and the solvent removed in vacuo. Chromatography on silica gel eluting with dichloromethane/methanol/35% ammonia solution (20:1:0.1) gave the title compound (0.12 g, 80%) as a colourless solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>) *inter alia* 0.81 (3H, d, J

6.5Hz), 1.08 (3H, d, J 7.1Hz), 1.15 (3H, s), 1.55 (3H, s), 2.26 (3H, s), 3.12 (2H, s), 5.20 (1H, dd, J 17.5 and 1.4Hz), 5.36 (1H, dd, J 10.9 and 1.4Hz), 5.72 (8.6Hz), 6.04 (1H, d, J 6.1Hz), 6.47 (1H, dd, J 17.5 and 10.9Hz), 7.73 (1H, d, J 6.1Hz); MS (+ve ion electrospray) 504 (MH<sup>+</sup>).

5

**Example 58 - Mutilin 14-(*exo*-8-methyl-8-azabicyclo[3.2.1]oct-3-ylsulfanyl)-acetate**  
22-Deoxy-22-sulfanylpleuromutilin (US Patent 4130709, 1978) (0.1 g, 0.00025 mole) in ethanol (4 ml) was treated with sodium methoxide (0.014 g, 0.0026 mole) and the resulting mixture stirred for 30 minutes. A solution of endo-3-methanesulfonyloxy-8-methyl-8-azabicyclo [3.2.1] octane (prepared from endo-8-methyl-8-azabicyclo [3.2.1] octan-3-ol and methanesulfonyl chloride) (0.061 g, 0.00028 mole) in ethanol (1 ml) was then added. Stirring was continued for 68 hours; a further portion of endo-3-methanesulfonyloxy-8-methyl-8-azabicyclo [3.2.1] octane (0.061 g, 0.00028 mole) was then added and stirring continued for a further 18 hours. The mixture was then diluted with dichloromethane, washed twice with aqueous potassium carbonate, once with brine, dried over magnesium sulfate and concentrated in vacuo. Chromatography on silica gel eluting with chloroform/methanol/35% ammonia solution (9:1:0:1) gave the title compound 0.035 g (27%), identical to the compound described in Example 50.

**Example 59 - Mutilin 14-(1-carboxamidomethylpiperidin-4-ylsulfanyl)-acetate**  
Mutilin (1-carboxymethylpiperidin-4-ylsulfanyl)-acetate (Example 37) (0.08 g, 0.00015 mole) in dichloromethane (3 ml) was treated with oxalyl chloride (0.032 ml, 0.00036 moles) and dimethylformamide (1 drop) and stirred at ambient temperature for 2 hours. The mixture was then evaporated to dryness and the residue suspended in tetrahydrofuran (3 ml) and treated with 35% aqueous ammonia solution (25 ml) and stirred for 2 hours. The mixture was evaporated to dryness and the residue positioned between saturated sodium bicarbonate and chloroform. The organic layer was separated and dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to dryness. Chromatography was saturated on silica gel eluting with chloroform/methanol/35% aqueous ammonia solution (90:9:1). Trituration of the residue obtained with methanol/diethyl ether gave the title compound 0.035 g; M.S. (+ve ion electrospray) m/z 535(MH<sup>+</sup>, 88%).

#### Antibacterial Activity

The following Table illustrates the antibacterial activities of representative mutilin 14-esters. Activities are given as minimum inhibitory concentrations in micrograms per millilitre (10<sup>-6</sup> g/ml), and were determined using a standard broth dilution method in microtitre.



Organism	Pleuromutilin	Tiamulin	Compound from Example 1	Compound from Example 15	Compound from Example 50
S.a.	2	0.25	$\leq 0.06$	$\leq 0.06$	$\leq 0.06$
S.p.	8	0.25	$\leq 0.06$	$\leq 0.06$	$\leq 0.06$
E.c.	> 64	> 64	16	64	32
H.i.	2	2	0.25	0.5	0.5
M.c.	0.5	0.125	$\leq 0.06$	$\leq 0.06$	$\leq 0.06$

S.a. = *Staphylococcus aureus* Oxford; S.p. = *Streptococcus pneumoniae* 1629;

E.c. = *Escherichia coli* DC0; H.i. = *Haemophilus influenzae* Q1;

M.c. = *Moraxella catarrhalis* Ravasio

## Pharmaceutical Compositions

### Example 1 – Oily Spray Formulation

- 5 A carrier for a nasal spray formulation was prepared by forming a blend of 67% w/w fractionated coconut oil (medium chain length)\* and 33% w/w of glyceryl mono-oleate \*\*. To this blend was added 0.2% w/w of powdered lemon juice flavour, followed by 0.5 or 1.0% w/w of drug substance (either in solution or, if insoluble, micronized)\*\*\*.
- 10 The resultant formulation has a viscosity which is sprayable at 20°C or above. When sprayed into the nose of a patient, the liquid coats the nasal passages and contact with moisture inside the nose (from the mucous membranes, and the humid environment generally) causes the carrier to thicken. This prolongs the residence time of the sprayed formulation on the nasal surfaces. A spray volume of about 100 µl contains approximately
- 15 0.5 or 1mg of drug substance.

\*Commercial product Miglyol, obtainable from Condea.

\*\* Commercial product Myverol 18-99, obtainable from Eastman.

\*\*\* for example, the compound of Example 1 or Example 8.

20



### Example 2 – Aqueous Spray formulation

	Component	%	Purpose
25	Drug	0.001-1.00	Active
	Sodium Chloride	0.5-0.9	Tonicity modifier
	Benzalkonium chloride	0.02	Preservative
	Disodium edetate	0.1	Part of preservative system/chelating agent
30	Polysorbate 80	0.2	Surfactant/solubiliser
	Sodium dihydrogen orthophosphate	0.2	Buffer
	Water	qs	Carrier

- Hydrochloric acid and sodium hydroxide were used to adjust the pH of the composition to
- 35 about pH 5.5. The drug molecule shows optimum stability at this pH.

## Claims

1. A compound of general formula (IA) or (IB):



in which:

- 10 each of n and m is independently 0, 1 or 2;  
 X is selected from -O-, -S-, -S(O)-, -SO<sub>2</sub>-, -CO.O-, -NH-, -CONH-, -NHCONH- and a bond;  
 R<sup>1</sup> is vinyl or ethyl;  
 R<sup>2</sup> is a non-aromatic monocyclic or bicyclic group containing one or two basic nitrogen  
 15 atoms and attached through a ring carbon atom;  
 R<sup>3</sup> is H or OH; or  
 the moiety R<sup>2</sup>(CH<sub>2</sub>)<sub>m</sub>X(CH<sub>2</sub>)<sub>n</sub>CH<sub>2</sub>COO at position 14 of (IA) or (IB) is replaced by  
 R<sup>a</sup>R<sup>b</sup>C=CHCOO in which one of R<sup>a</sup> and R<sup>b</sup> is hydrogen and the other is R<sup>2</sup> or R<sup>a</sup> and R<sup>b</sup>  
 together form R<sup>2</sup>; or  
 20 a pharmaceutically acceptable salt thereof.
2. A compound according to claim 1 in which R<sup>2</sup> is selected from optionally  
 substituted piperidinyl, pyrrolidyl, quinuclidinyl, azabicyclo[2.2.1]heptyl,  
 azabicyclo[4.3.0]nonyl, azabicyclo[3.2.1]octyl, azabicyclo[3.3.0]octyl,  
 25 azabicyclo[2.2.2]octyl, azabicyclo[3.2.1]octenyl, azabicyclo[3.3.1]nonyl and  
 azabicyclo[4.4.0]decyl.

3. A compound according to claim 1 or 2 in which R<sup>2</sup> is substituted by alkyl, alkyloxy, alkenyl or alkenyloxy, which are optionally further substituted by one or more groups selected from aryl, heterocyclyl, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)alkylthio, aryl(C<sub>1-6</sub>)alkoxy, aryl(C<sub>1-6</sub>)alkylthio, amino, mono- or di-(C<sub>1-6</sub>)alkylamino, cycloalkyl, cycloalkenyl, carboxy and esters thereof, amides of carboxy, ureido, carbamimidoyl (amidino), guanidino, alkyl-sulfonyl, amino-sulfonyl (C<sub>1-6</sub>)acyloxy, (C<sub>1-6</sub>)acylamino, azido, hydroxy, and halogen.
4. A compound according to claim 1 selected from
- mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate
  - mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate
  - 19,20-dihydromutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate
  - mutilin 14-(quinuclidin-3-yloxy)-acetate
  - mutilin 14-(quinuclidin-3-ylsulfanyl)-acetate
  - mutilin 14-(quinuclidin-4-yl-sulfanyl)-acetate
  - mutilin 14-[N-(2,2-dimethylazabicyclo[4.3.0]non-4-ylmethyl)]-aminoacetate
  - mutilin 14-(quinuclidin-4-ylcarbonylamino)-acetate
  - mutilin 14-[(3R,4R)-azabicyclo[2.2.1]hept-3-ylcarbonylamino]-acetate
  - mutilin 14-(1-methylpiperid-4-ylcarbonylamino)-acetate
  - mutilin 14-[3-(1-methylpiperid-4-yl)]-propionate
  - mutilin 14-(quinuclid-4-ylmethysulfanyl)-acetate
  - 19,20-dihydromutilin 14-(quinuclidin-4-ylsulfonyl)acetate
  - 19,20-dihydromutilin 14-(quinuclidin-4-ylsulfoxy)-acetate
  - mutilin-14-(1-methylpiperid-4-ylsulfanyl)-acetate
  - mutilin 14-[(3RS,4SR)-1-aza-bicyclo[2.2.1]hept-3-yl-sulfanyl]-acetate
  - mutilin 14-[(3RS,4SR)-1-aza-bicyclo[2.2.1]hept-3-yl-sulfanyl]-acetate
  - mutilin 14-(quinuclidin-3-ylidene)-acetate
  - mutilin 14-[quinuclidin-3-yl]-acetate
  - mutilin 14-[quinuclidin-3-ylacetoxyl]-acetate
  - mutilin 14-(quinuclidin-3-ylmethysulfanyl)-acetate
  - 1,2-didehydromutilin 14-(quinuclidin-4-ylsulfanyl)-acetate
  - 2 $\alpha$ -hydroxymutilin 14-(quinuclidin-4-ylsulfanyl)-acetate
  - mutilin 14-(quinuclidin-4-yl)-acetate
  - mutilin 14-(quinuclidin-4-ylmethyl)-aminoacetate
  - mutilin 14-[3-(quinuclidin-4-yl)-acrylate]
  - mutilin 14-[3-(quinuclidin-4-yl)]-propionate
  - mutilin 14-(quinuclidin-4-ylmethyloxy)-acetate

- mutilin 14-[(3R)-quinuclidin-3-ylamino]-acetate  
 mutilin 14-(quinuclidin-4-yl-amino)-acetate  
 mutilin 14-[4-(quinuclidin-4-yl)]-butyrate  
 mutilin 14-(1-azabicyclo[3,3,0]oct-4-ylmethylsulfanyl)-acetate  
 5 mutilin 14-(1-azabicyclo[3,3,0]oct-3-ylsulfanyl)-acetate  
 mutilin 14-(endo 8-methyl-8-azabicyclo[3,2,1]oct-3-ylsulfanyl)-acetate  
 mutilin 14-(1-azabicyclo[4,3,0]non-4-ylsulfanyl)-acetate  
 19,20-dihydromutilin 14-(1-azabicyclo[4,3,0]non-4-ylsulfanyl)-acetate  
 mutilin 14-(1-carboxymethylpiperidin-4-ylsulfanyl)-acetate  
 10 mutilin 14-(piperidin-4-ylsulfanyl)-acetate  
 mutilin 14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate  
 mutilin 14-[(3S,4R)-1-azabicyclo[2.2.1]hept-3-ylmethylsulfanyl]-acetate  
 mutilin 14-(quinuclidin-2-ylmethylsulfanyl)-acetate  
 mutilin 14-(1-azabicyclo[2.2.1]hept-4-ylmethylsulfanyl)-acetate  
 15 mutilin 14-[(3R,4S)-1-azabicyclo[2.2.1]hept-3-ylmethylsulfanyl]-acetate  
 mutilin 14-(1-azabicyclo[3.2.1]oct-5-ylmethylsulfanyl)-acetate  
 mutilin 14-[(R)-1-methylpiperid-2-ylmethylsulfanyl]-acetate  
 mutilin 14-[(S)-1-methylpyrrolid-2-ylmethylsulfanyl]-acetate  
 mutilin 14-[(R)-1-methylpiperid-3-ylmethylsulfanyl]-acetate  
 20 mutilin 14-(quinuclidin-4-ylmethylsulfanyl)-acetate  
 mutilin 14-(8-methyl-8-azabicyclo[3.2.1]oct-3-ylmethylsulfanyl)-acetate  
 mutilin 14-(exo-8-methyl-8-azabicyclo[3.2.1]oct-3-ylsulfanyl)-acetate  
 mutilin 14-[(3-(quinuclidin-4-ylsulfanyl))-propionate  
 mutilin 14-[3-(quinuclidin-4-ylmethylsulfanyl)]-propionate  
 25 mutilin 14-[3-(1-methylpiperid-4-ylsulfanyl)]-propionate  
 19,20-dihydromutilin 14-(1-methylpiperid-4-ylsulfanyl)-acetate  
 19,20-dihydromutilin 14-(8-methyl-8-azabicyclo[3.2.1]oct-3-ylmethylsulfanyl)-acetate  
 mutilin 14-[4-(quinuclidin-4-ylsulfanyl)]-butyrate  
 1,2-didehydromutilin 14-(1-methylpiperidin-4-ylmethylsulfanyl)-acetate  
 30 mutilin 14-(exo-8-methyl-8-azabicyclo[3.2.1]oct-3-ylsulfanyl)-acetate, or  
 mutilin 14-(1-carboxamidomethylpiperidin-4-ylsulfanyl)-acetate.

5. A process for preparing a compound according to any one of claims 1 to 4 which comprises:
- 35 (a) coupling mutilin or epi-mutilin having a protected hydroxy group at position 11, with an active derivative, such as an acid chloride, of a carboxylic acid  
 $R^{2A}-(CH_2)_m-X-(CH_2)_n-CH_2CO_2H$ , where  $R^{2A}$  is  $R^2$  as defined in claim 1 or a group convertible thereto, and n, m, and X are as defined in claim 1, and if necessary converting

the epi-mutilin to mutilin, and where necessary or desired, before or after the coupling, modifying the mutilin nucleus to introduce 2-OH; 19,20-dihydro; or 1,2-dehydro substituents; or

- 5 (b) providing a mutilin or epi-mutilin derivative having  $(\text{CH}_2)_n\text{CH}_2\text{CO}$  as an *O*-acyl group at position 14, where the acyl group is substituted with  $\text{R}^{\text{L}}$ , which is a leaving group, OH or NH, coupling the 14-*O*-acyl-(epi)mutilin derivative with a compound  $\text{R}^2\text{A}(\text{CH}_2)_m\text{XH}$  or an active derivative thereof, and if necessary converting the epi-mutilin configuration to mutilin, and where necessary or desired, before or after the coupling, modifying the mutilin  
10 nucleus to introduce 2-OH; 19,20-dihydro; or 1,2-dehydro substituents.

6. A process for preparing a compound according to claim 5(b) in which

(a) when X is O, S or NH,  $\text{R}^{\text{L}}$  is a leaving group and is reacted with

- 15 (i) the alcohol  $\text{R}^2-(\text{CH}_2)_m\text{-OH}$ ;  
(ii) the thiol  $\text{R}^2-(\text{CH}_2)_m\text{-SH}$ ;  
(iii) the amine  $\text{R}^2-(\text{CH}_2)_m\text{-NH}_2$ ;

(b) when X is CONH,  $\text{R}^{\text{L}}$  is amino and is reacted with the acid  $\text{R}^2\text{A}-(\text{CH}_2)_m\text{-CO}_2\text{H}$ ,  
or an acylating agent derived therefrom;

- 20 (c) when X is CO.O,  $\text{R}^{\text{L}}$  is hydroxy and is reacted with an acylating agent derived from the acid  $\text{R}^2\text{A}-(\text{CH}_2)_m\text{-CO}_2\text{H}$ .


7. A pharmaceutical composition comprising a compound according to claim 1 and a  
25 pharmaceutically acceptable carrier.

8. A pharmaceutical composition according to claim 7 in the form of a spray adapted for administration to the nasal cavity.

30 9. A pharmaceutical composition according to claim 8 in which the spray is an aqueous spray.

10. A compound of formula (IA) or (IB) according to claim 1 for use in therapy.

11. The use of a compound of formula (IA) or (IB) according to claim 1 in the manufacture of a medicament adapted for administration to the nasal cavity for reducing or eliminating the nasal carriage of pathogenic organisms.
- 5
12. The use of a compound of formula (IA) or (IB) according to claim 1 in the manufacture of a medicament adapted for administration to the nasal cavity for prophylaxis of recurrent otitis media or recurrent acute bacterial sinusitis.
- 10
13. The use of a compound of formula (IA) or (IB) according to claim 1 in the manufacture of a medicament for treating skin and soft tissue infections and acne.
14. A method of treating microbial infections in animals, especially in humans and in domesticated mammals, which comprises administering a compound according to claim 1, to a patient in need thereof.
- 15
15. A method of reducing or eliminating the nasal carriage of pathogenic organisms which method comprises administering to the nasal cavity a compound a compound according to claim 1, to a patient in need thereof.
- 20
16. A method of prophylaxis of recurrent otitis media or recurrent acute bacterial sinusitis in humans , which comprises administering to the nasal cavity a compound according to claim 1, to a patient in need thereof.
- 25
17. A method of treatment of skin and soft tissue infections and acne in humans , which comprises topically administering a compound according to claim 1, to a patient in need thereof.
18. A compound of the general formula (I):



in which:

each of n and m is independently between 0 and 2;

X is selected from the group consisting of -O-, -S-, -S(O)-, -SO<sub>2</sub>-, -NH-, CONH-, -CH<sub>2</sub>-,

5 and a bond;

R<sup>1</sup> is vinyl or ethyl; and

R<sup>2</sup> is a non-aromatic monocyclic or bicyclic group containing one or two basic nitrogen atoms and attached through a ring carbon atom.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03211

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D453/02 A61K31/435 C07D451/02 C07D471/08 C07D487/08  
C07D211/62 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 98 05659 A (NAYLOR ANTOINETTE ;HUNT ERIC (GB); SMITHKLINE BEECHAM PLC (GB); TA) 12 February 1998 cited in the application see claim 1	1-18
X	WO 97 25309 A (SMITHKLINE BEECHAM PLC ;HINKS JEREMY DAVID (GB); TAKLE ANDREW KENN) 17 July 1997 cited in the application see page 4, line 15 - line 22; claim 1	1-18
X	DE 34 05 632 A (SANDOZ AG) 22 August 1985 see claim 1; examples 7,8,11	1-18
X	US 4 278 674 A (EGGER HELMUT ET AL) 14 July 1981 see claim 1	1-18
	--- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

22 January 1999

Date of mailing of the international search report

05.02.99

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03211

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 013 768 A (SANDOZ AG) 6 August 1980 Examples 21.29,33 see page 5, line 7 - line 26; claim 1; examples 1-6,13,14,19 ---	1-18
X	GB 2 025 930 A (SANDOZ LTD) 30 January 1980 see claim 1 ---	1-18
X	US 4 060 542 A (RIEDL KURT) 29 November 1977 see claim 1 ---	1-18
X	EGGER H ET AL: "NEW PLEUROMUTILIN DERIVATIVES WITH ENHANCED ANTIMICROBIAL ACTIVITY.II. STRUCTURE-ACTIVITY CORRELATIONS" JOURNAL OF ANTIBIOTICS, vol. 29, no. 9, September 1976, pages 923-927, XP002028938 cited in the application see the whole document -----	1-18

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/GB 98/03211

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 13-17 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

There is a contradiction between the independent claim 1 and the dependent claim 2 since claim 1 does not allow for optional substitution of the R2 group whereas claim 2 does. It is assumed that this is a mistake and the search has been carried out on the basis that R2 in claim 1 is substituted as described on pages 3-4 of the description.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03211

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9805659 A	12-02-1998	AU 1307897 A AU 4203697 A EP 0874809 A NO 983074 A CA 2240467 A WO 9725309 A	01-08-1997 25-02-1998 04-11-1998 31-08-1998 17-07-1997 17-07-1997
WO 9725309 A	17-07-1997	AU 1307897 A CA 2240467 A EP 0874809 A NO 983074 A AU 4203697 A WO 9805659 A	01-08-1997 17-07-1997 04-11-1998 31-08-1998 25-02-1998 12-02-1998
DE 3405632 A	22-08-1985	AT 29132 T AU 585209 B AU 3877385 A CA 1249281 A CY 1565 A DD 253616 A DK 72885 A,B, EP 0153277 A FI 850654 A,B, FI 921848 A,B, GR 850415 A HK 103690 A IE 58232 B JP 1886041 C JP 6008278 B JP 60199870 A LV 5764 A SU 1582985 A US 4675330 A	15-09-1987 15-06-1989 12-09-1985 24-01-1989 17-05-1991 27-01-1988 18-08-1985 28-08-1985 18-08-1985 24-04-1992 18-06-1985 21-12-1990 11-08-1993 22-11-1994 02-02-1994 09-10-1985 20-12-1996 30-07-1990 23-06-1987
US 4278674 A	14-07-1981	CH 572893 A CH 568272 A CH 575375 A CH 560180 A CH 559719 A AT 335054 B AT 845872 A BE 789629 A BG 21386 A BG 20773 A CA 979447 A CS 219306 B CY 987 A DD 100706 A DE 2248237 A DE 2265666 C DK 139720 B DK 498076 A,B, DK 498176 A,B, FI 58635 B FI 794060 A,B, FI 794061 A,B, FR 2157828 A GB 1410506 A	27-02-1976 31-10-1975 14-05-1976 27-03-1975 14-03-1975 25-02-1977 15-06-1976 03-04-1973 20-05-1976 20-12-1975 09-12-1975 25-03-1983 02-08-1979 05-10-1973 12-04-1973 03-02-1983 02-04-1979 03-11-1976 03-11-1976 28-11-1980 27-12-1979 27-12-1979 08-06-1973 15-10-1975

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03211

Patent document cited in search report		Publication date	Patent family member(s)	Publication date		
US 4278674	A		GB 1410505 A	15-10-1975		
			HK 25379 A	27-04-1979		
			IE 37699 B	28-09-1977		
			IE 37700 B	28-09-1977		
			JP 48044244 A	26-06-1973		
			NL 7213254 A,B,	09-04-1973		
			OA 4192 A	31-12-1979		
			SE 398348 B	19-12-1977		
			SE 424074 B	28-06-1982		
			SE 7509456 A	26-08-1975		
			SE 419984 B	07-09-1981		
			SE 7509457 A	26-08-1975		
			US 4107164 A	15-08-1978		
			US 4208326 A	17-06-1980		
			AT 335055 B	25-02-1977		
			AT 735974 A	15-06-1976		
			CS 219307 B	25-03-1983		
			US 4032530 A	28-06-1977		
			AT 341666 B	27-02-1978		
			AT 736074 A	15-06-1977		
			CS 219308 B	25-03-1983		
			AU 4740572 A	11-04-1974		
			JP 56095163 A	01-08-1981		
			JP 1219330 C	26-07-1984		
			JP 56095180 A	01-08-1981		
			JP 58052994 B	26-11-1983		
		-----				
		EP 0013768	A	06-08-1980	AT 6062 T	15-02-1984
					AU 533691 B	08-12-1983
					AU 5453580 A	17-07-1980
CA 1125778 A	15-06-1982					
CY 1353 A	24-04-1987					
DK 4280 A,B,	13-07-1980					
FI 800024 A,B,	13-07-1980					
HK 40587 A	29-05-1987					
IE 49356 B	18-09-1985					
JP 55094359 A	17-07-1980					
KE 3689 A	20-02-1987					
US 4428953 A	31-01-1984					
ZA 8000175 A	26-08-1981					
-----						
GB 2025930	A	30-01-1980	NONE			
-----						
US 4060542	A	29-11-1977	AT 301753 B	15-08-1972		
			US 3979423 A	07-09-1976		
			CA 986102 A	23-03-1976		
			CA 970387 A	01-07-1975		
			CS 172913 B	28-01-1977		
			DE 2036027 A	18-02-1971		
			DE 2066118 B	28-08-1980		
			DK 66077 A	16-02-1977		
			FR 2059559 A	04-06-1971		
			GB 1312148 A	04-04-1973		
			NL 7010444 A	27-01-1971		
			SE 369712 B	16-09-1976		
			ZA 7005091 A	23-02-1972		
-----						